

Faktor-faktor yang berhubungan dengan Indeks Massa Tubuh (IMT) remaja putri di Kota Bukittinggi Tahun 2006 = Factors related to the Body Mass Index (BMI) of girls in Bukittinggi in 2006

Rini Santy, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=106943&lokasi=lokal>

Abstrak

Kejadian gizi kurang pada remaja putri (rematri) sering terlupakan dari penglihatan dan pengamatan biasa, padahal kualitas sumber daya manusia (SDM) sebagai indikator keberhasilan pembangunan nasional terletak ditangan remaja. Menurut Susenas 1999-2003, 35 - 40% Wanita Usia Subur (WUS) 15-19 tahun berisiko Kekurangan Energi Kronis (ICED). Keadaan gizi kurang merupakan akibat dari asupan energi yang tidak cukup. Salah satu cara untuk menentukan keadaan gizi seseorang adalah dengan mengetahui Indeks Massa Tubuh (IMT) yaitu membandingkan berat badan dan tinggi badan (kg/m^2).

Penelitian ini bertujuan mendapatkan gambaran status gizi remaja putri di Kota Bukittinggi dan faktor-faktor yang berhubungan. Penelitian ini merupakan analisis data primer dengan pendekatan kuantitatif observasional. Rancangan penelitian adalah potong lintang (cross sectional). Penelitian dilaksanakan pada bulan Februari-Maret 2006. Remaja putri pada penelitian ini diwakili oleh siswi kelas III SLTA (SMA, MA, dan SMK) usia 16 - 18 tahun yang dikategorikan remaja akhir yang sangat dekat dengan masa kehamilan. Pemilihan sampel dilakukan secara systematic random sampling. Sampel berjumlah 156 orang yang tersebar pada 11 sekolah. Variabel terikat adalah IMT dan variabel bebas adalah asupan energi, kebiasaan makan, citra tubuh, pengetahuan gizi, kelompok sebaya, aktivitas fisik, dan karakteristik orang tua. Analisis data dilakukan secara bertahap dimulai dengan univariat, bivariat (chi square) dan multivariat (multiple logistic regression).

Hasil penelitian menunjukkan rata-rata IMT rematri adalah $20,69 \text{ kg}/\text{m}^2 + 2,63$. Proporsi siswi yang mempunyai $\text{IMT} < 18,5 \text{ kg}/\text{m}^2$ sebesar 19,9% dengan penyebaran 14,1% kekurangan gizi tingkat ringan dan 5,8% kekurangan gizi tingkat berat. Rata-rata asupan energi adalah 1694 kalori. Rata-rata kontribusi protein terhadap total energi sebesar 11,8%, lemak 26,7% dan karbohidrat 58,7%. Rata-rata asupan energi dibandingkan AKG adalah total energi 77%, protein 93,6%, lemak 65,3% dan karbohidrat 84,7%.

Hasil analisis bivariat menunjukkan ada hubungan bermakna antara total energi, kebiasaan makan dan citra tubuh dengan IMT rematri. Variabel total energi merupakan variabel yang dominan mempengaruhi status gizi IMT rematri.

Berdasarkan hasil penelitian, disarankan kepada pengambil keputusan bidang kesehatan agar menyusun program penanggulangan dan pencegahan masalah gizi remaja. Kegiatan yang dapat dilakukan antara lain aryaia mengadakan pelatihan untuk petugas khusus promosi gizi dan pelatihan guru BP/guru olah raga mengenai pemantauan status gizi, melaksanakan promosi gizi secara intensif dengan lebih mengarahkan sasaran penyuluhan ke sekolah serta membuat sarana penyuluhan yang disesuaikan dengan karakter remaja. Untuk Dinas Pendidikan agar dapat mengintegrasikan materi kesehatan khususnya pengetahuan gizi

ke dalam kurikulum, menggiatkan UKS dan KKR untuk mengoptimalkan penggunaan KMS anak sekolah, menyediakan ruang UKS yang dilengkapi dengan timbangan, microtoise, food model dan buku-buku gizi, melaksanakan PSG setiap awal semester dan bekerja sama dengan orang tua siswa untuk dapat menyediakan makan siang di sekolah (school lunch) guna menjaga asupan yang adekuat mengingat sebagian besar waktu dihabiskan di sekolah. Untuk peneliti yang berminat melakukan penelitian status gizi remaja agar menggunakan indikator status gizi yang memperhitungkan pacu tumbuh yang sesuai dengan remaja Indonesia serta penelitian citra tubuh secara mendalam yang diduga mempengaruhi perilaku makan remaja.

<hr><i>The incident of malnutrition at girls is often neglected from common sight and observation, whereas the quality of human resources as the indicator of a successful national development is laid on their hand. According to the National Health Survey (Susehns) of 1999 2003, 35 - 40% women in productive age (WUS) of 15 - 19 are at risk of Chronic Energy Deficiency (KEK). Malnutrition is resulted from the insufficient consumption of energy. One of ways to determine the nutritional condition of a person is finding out the Body Mass Index (BMI) of him/her, namely by comparing his/her body weight with his/her height (kg/mi).

This research is aimed at obtaining the description of nutritional status of girls in Bukittinggi and factors related to it. It was conducted by analyzing primary data using observational quantitative approach. The design of the research is cross sectional. The research was carried out from February until March 2006. The girls studied are represented by the third-grade female students of senior high schools (senior high school, Islamic senior high school, and middle vocational school) of 16 -18 who are categorized as a last teenager who is very close to pregnant period. The sample was selected by systematic random sampling. It was totally 156 students who are distributed at 11 schools. The dependent variable is BMI and the independent variable are energy consumption, eating habit, body image, knowledge on nutrition, peer group, physical activities, and parents' characteristics. Data was analyzed gradually, starting from univariate, bivariate (chi square), until multivariate (multiple logistic regression).

The results show that the BMI of the female students is $20.69 \text{ kg/m}^2 \pm 2.63$ on average. The proportion of students having $\text{BMI} < 18.5 \text{ kg/m}^2$ is 19.9% all of which is distributed to 14.1% of light level of malnutrition and 5.8% for heavy level of malnutrition. Intake per day is 1694 calorie on average with protein contributed to intake is 11,8%, fat 26,7% and carbohydrate 58,7%. Intake energy compared with Recommended Dietary Allowance (RDA) are total energy consumption 77%, protein 93,6%, lemak 65,3% and carbohydrate 84,7%.

Bivariate analysis indicates that there is a significant relation between energy consumption, eating habit, body image, by BMI. Variable energy consumption the dominant variable influencing BMI.

Based on the results, it is suggested that the decision maker in health areas begin to set up prevention and control program for nutritional problems of teenagers. Activities which can be conducted among others are training for special personnel of nutritional promotion and BP/sport teachers on nutritional status monitoring, conducting nutritional promotion intensively which is more focusing on education at schools, and setting up educational facilities adjusted to teenager character. It is also recommended that the Educational Office integrate health matters, especially nutritional knowledge into the curriculum, activate UKS and KKR to optimize the using of KMS of school students, provide UKS room equipped with scale,

microtoise, food model and nutrition books, conduct PSG at the beginning of every semester and cooperate with students' parents to provide school lunch to maintain adequate intake, considering that most of their time is spent at school. Suggestion for the researcher to use nutritional status that adjusted growth spurts Indonesian girls and study of factor body image which influence food habit.</i>