

Analisis faktor risiko yang berpengaruh terhadap skor kardiovaskular Jakarta dan skor Framingham pada pekerja PT-X Jakarta, 2004 =
Analysis of the risk factors that influence Jakarta cardiovascular score and Framingham score among PT-X employees in Jakarta, 2004

Ida Bagus Adiatmaja, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=107024&lokasi=lokal>

Abstrak

Latar belakang

Penyakit kardiovaskular merupakan penyebab kematian utama pada pekerja perusahaan yang bergerak dalam minyak dan gas bumi nasional. Para pekerja tersebut diharapkan mempunyai kewaspadaan akan faktor risiko penyakit kardiovaskular tersebut.

Tujuan penelitian ini adalah untuk mengetahui apakah Skor Kardiovaskular Jakarta dapat dipakai guna menentukan kemungkinan kejadian kardiovaskular, guna upaya promotifpreventif risiko dan mengetahui hubungan faktor pekerjaan dengan Skor Kardiovaskular Jakarta maupun Skor Framingham.

Metode

Penelitian ini menggunakan rancangan kros sektional dengan 107 responden. Dilakukan pengumpulan data primer melalui wawancara, pengisian kuesioner dan pengumpulan data sekunder melalui penelusuran catatan rekam medic.

Hasil

Karakteristik sosiodemografi subyek sebagian besar berumur > 45 tahun (70,1%). Skor Kardiovaskular Jakarta subyek sebagian besar tergolong kategori risiko tinggi (58,0%), sedangkan Skor Framingham subyek sebagian besar risiko rendah (76,6%). Dari analisis regresi logistik binary yang paling kuat berhubungan dengan Skor Kardiovaskular Jakarta adalah umur (OR-suaian=10,06, 95% CI=2,43-41,66), sedangkan yang paling kuat berhubungan dengan Skor Framingham adalah diabetes melitus (OR-suaian=216,82, 95% CI=13,76-3416,07) dan kolesterol-total (OR-suaian=162,17, 95% CI=13,27-1982,17). Terdapat korelasi yang bermakna dengan arah positif dan cukup kuat antara Skor Kardiovaskular Jakarta dengan Skor Framingham (koefisien korelasi = 0,592 dan p = 0,000).

Kesimpulan

Skor Kardiovaskular subyek sebagian tergolong kategori risiko tinggi. Skor Framingham subyek sebagian besar tergolong kategori rendah. Tidak ada hubungan faktor pekerjaan dengan kedua skor tersebut. Terdapat korelasi antara Skor Kardiovaskular Jakarta dengan Skor Framingham. Skor Kardiovaskular Jakarta dapat dipakai pada populasi penelitian ini.

<hr><i>Background

Cardiovascular diseases are among the most common causes of death in employees of the national oil and gas company. Employees should be made aware on the cardiovascular risk factors.

The aim of this research was to know if Jakarta Cardiovascular Score could be used to determine cardiovascular risks and to know the relationship between job factors, Jakarta Cardiovascular Jakarta and Framingham Score.

Methods

This study was using cross sectional design with a sample of 107 respondents. Data were collected by interview using questionnaire and medical record file review.

Result

Sociodemographic characteristics of the respondents showed that most of them were 45 years of age. The study found out that using Jakarta Cardiovascular Score most subjects showed risk high (58,0%), while using Framingham Score most subjects still showed risk low (76,6%). The result of logistic binary regression indicated that there were significant relationship between age and Jakarta Cardiovascular Score (adjusted-OR= 10,06, 95% CI=2,43-41,66) and also there were significant relationship among diabetic, cholesterol level and Framingham Score (adjusted-OR--216,82, 95% CI-13,76-3416,09 and adjusted-CR=162,17, 95% CI=13,27-1982,17). No significant relationship was found between job factors and either scores. A positive significant correlation was established between Jakarta Cardiovascular Score and Framingham score (coefficient of correlation 0,592, $p=0,000$).

Conclusion

Most subject showed high Jakarta Cardiovascular Score, while most subject showed low Framingham Score. No significant relationship was found between job factors and either scores. A positive significant correlation was established between Jakarta Cardiovascular Score and Framingham Score. Jakarta Cardiovascular Score can be used for the populations of this study.