

Intestinal parasitic infections in primary school children in Pulau Panggang and Pulau Pramuka, Kepulauan Seribu / Adi Sasongko, Heksa SJY Irawan, Rahmi S Tatang, Riza Subahar, Purnomo, Sri S Margono

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=117241&lokasi=lokal>

Abstrak

Pada survei ini telah dikumpulkan sejumlah sampel tinja untuk pemeriksaan terhadap cacing yang ditularkan melalui tanah dan terhadap Protozoa pada anak kelas tiga sekolah dasar. Pemeriksaan tersebut merupakan bagian program penanggulangan cacing yang ditularkan melalui tanah. Ketiga sekolah berlokasi di dua pulau yaitu Pulau Panggang dan Pulau Pramuka, yang termasuk sekelompok pulau dekat pantai Utara Jakarta. Sekolah-sekolah ini belum pernah ikut serta dengan kegiatan program penanggulangan S-TH. Untuk pemeriksaan digunakan cara semi kuantitatif Kato sediaan tebal dan cara sediaan langsung dengan larutan yodium 2%. Empat spesies cacing dan lima spesies Protozoa telah ditemukan pada 101 sampel tinja. Infeksi Ascaris dan Tichuris ditemukan sebanyak 68.8% atau lebih. Infeksi cacing tambang hanya ditemukan di satu sekolah (2.9%). Telur *Hymenolepis nana* ditemukan pada satu sampel. Diantara 101 sample ini ditemukan 5% *Entamoeba histolytica* dan *Entamoeba coli*, sedangkan 2% *Endolimax nana* ditemukan pada 2.0% diantara sampel tinja ini. Tercatat prevalensi tinggi untuk *Blastocystis hominis* (36.0%) dan *Giardia lamblia* (30.0%). Kebanyakan infeksi Ascaris termasuk infeksi ringan di SD I (69.0%) dan tidak ditemukan infeksi berat di SD ini. Kebanyakan infeksi di SD II dan III merupakan infeksi sedang yaitu untuk masing-masing sekolah 51.4 dan 81.8%. Di SD II dan III juga ditemukan infeksi berat yaitu untuk masing-masing sekolah 11.4 dan 5.8%. Telur Ascaris yang dibuahi ditemukan pada 93.1%, 100% dan 95.5% berturut-turut di SD I, II dan III. Diantara 86 sampel yang positif 96.5% sampel mengandung telur yang dibuahi, sedangkan pada 3.5% sampel ditemukan telur yang tidak dibuahi. Prevalensi infeksi Ascaris dan Trichuris dapat diduga sebelumnya mengingat derajat hygiene lingkungan dan sanitasi yang rendah di daerah ini. Diantara infeksi Protozoa spesies yang dominan adalah *B. hominis* dan *G. lamblia*.

Stool samples were collected and examined for soil-transmitted helminthic and protozoal infection in the first grade of three primary schools, located on Pulau Panggang and Pulau Pramuka, which are parts of a group of islands not far from the north coast of Jakarta. The stool examinations were part of activities during a control program on soil-transmitted helminthic infections. The schools have never participated with control programs on soil-transmitted helminthiases. For the examination of the samples a semi-quantitative Kato thick smear method was used and the direct smear with a 2% iodine solution. Four intestinal helminth species and five protozoa species were found in a total of 101 stool samples. Ascaris and Trichuris infections were found in 68.8% or more. Hookworm infection was only found in one school (2.9%). Eggs of

Hymenolepis

nana were detected in one sample. Cysts of *Entamoeba histolytica* and *Entamoeba coli* were both found in 5.0% of the

samples, whereas *Endolimax nana* was recovered from 2.0% of the samples. High prevalence rates were detected for

Blastocystis hominis (36.0%) and for *Giardia lamblia* it was 30.0%. Most of the *Ascaris* infections were categorized as light

infections at School I (69.0%) and not a single heavy infection were found in this school. In School II and III most of the

infections were moderate i.e. respectively 51.4 and 81.8%. Also in Schools II and III heavy infections were detected,

respectively 11.4 and 5.8%. Fertilized *Ascaris* eggs were detected in 93.1%, 100% and 95.5% at School I, II and III respectively.

As a whole among 86 positive samples 96.5% were recorded as samples with fertilized eggs, whereas 3.5% contained

unfertilized eggs. The high prevalences of *Ascaris* and *Trichuris* infections in this area could be expected due to the low

level of environmental hygiene and sanitation. Among the protozoal infections *B. hominis* and *G. lamblia*

were the

dominant species.