

Perbedaan pH lambung pada pasien dispepsia dengan atau tanpa diabetes melitus tipe 2 = Differences gastric acid ph in dispepsia patients with or without dm type 2

Ilum Anam, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20365155&lokasi=lokal>

Abstrak

Latar Belakang: Sindroma dispepsia sering dialami oleh penderita DM. Asam lambung salah satu faktor agresif terjadinya sindroma dispepsia dan tukak lambung. Penelitian ini bertujuan untuk mencari perbedaan pH lambung pada pasien dispepsia DM dengan yang bukan DM dan untuk mengetahui apakah ada korelasi antara pH lambung dengan proteinuria dan HbA1c.

Metode: Pasien terdiri dari 30 kelompok DM dan 30 kelompok bukan DM. Masing-masing kelompok dihitung pH lambung basal. pH lambung basal diukur dgn memasukkan elektroda kateter kedalam lambung selama 30 menit kemudian di rekam dgn alat PH Metri merek Digitrapper pH-Z. Beratnya komplikasi DM diukur dengan mikroalbuminuria, sedangkan kendali gula darah diukur dgn HbA1c. Dilakukan uji chi square utk mencari perbedaan pH lambung kelompok DM dgn yg bukan DM, dengan terlebih dahulu menentukan titik potong dgn analisa ROC (Receiver Operating Characteristic). Dilakukan uji korelasi antara pH lambung basal dengan mikroalbuminuria dan HbA1c pada kelompok pasien DM.

Hasil: pH lambung basal pada dispepsia DM vs non DM (2.30 ± 0.83 vs 2.19 ± 0.52). Dgn uji chi square terdapat perbedaan bermakna antara kelompok DM dengan yang bukan DM. Pada uji korelasi antara pH lambung dengan mikroalbuminuria dijumpai $r = 0.47$ dan $p < 0.05$, sedangkan HbA1c dijumpai $r = 0.59$ dan $p > 0.05$.

Simpulan: Ada perbedaan bermakna pH lambung basal antara pasien dispepsia DM dengan pasien dispepsia bukan DM. Ada korelasi antara pH lambung basal dengan mikroalbuminuria, sedangkan dengan HbA1c tidak ada korelasi. pH lambung basal pada pasien DM adalah 2.03 ± 0.83 sedangkan pada yang bukan DM adalah 2.19 ± 0.52 .

.....

Aims: Dyspepsia syndrome often experienced in diabetic patients. Gastric acid was one aggressive factors in dyspepsia syndrome. This aim of this study was to determine differences gastric pH between dyspepsia diabetic and dispepsia without diabetic patients. Also to determine whether there were a correlation between basal gastric pH and microalbuminuria and also HbA1c.

Methods: There were 30 patients diabetic and 30 patients without diabetic. Basal gastric pH was measured with an electrode catheter that inserted into the stomach for 30 minutes. Gastric pH will be recorded with PH Metri Digitrapper pH-Z. Diabetic complications measured by microalbuminuria, while the measured blood sugar control with HbA1c. Chi-square test to determine differences gastric pH between diabetic and without diabetic patients. Correlation test was performed between basal gastric pH and microalbuminuria and also HbA1c.

Results: We found basal gastric pH diabetic and non diabetic patients were (2.30 ± 0.83 vs 2.19 ± 0.52). There was significant differences between diabetic and non diabetic patients. From 30 diabetic patients we found a corelation between basal gastric pH and microalbuminuria ($p < 0.05$ and $r = 0.47$) and a no corelation with HbA1c ($p > 0.05$ and $r = 0.59$).

Conclusions: There was significant differences basal gastric pH between diabetic and non diabetic patients. There was correlation between basal gastric pH and microalbuminuria, and no correlation with HbA1c. Basal gastric pH diabetic patients was $2,30 \pm 0,83$ and non diabetic patients was $2,19 \pm 0,52$.