

Uji sensitivitas dan spesifisitas Rapid Plasma Reagin dibandingkan dengan treponema pallidum haemagglutination assay sebagai baku emas diagnosis serologis frambusia pada anak usia 1,5 tahun di Kabupaten Sumba Barat Daya, Nusa Tenggara Timur periode 11-17 Juli 2010 = Sensitivity and specificity of Rapid Plasma Reagin compared to treponema pallidum haemagglutination assay as gold standard in serological diagnostic tool for yaws in children 1,5 years old at Sumba Barat Daya District, Nusa Tenggara Timur Province period 11-17 July 2010

Aruan, Rompu Roger, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20367215&lokasi=lokal>

---

Abstrak

Latar Belakang : Frambusia adalah infeksi yang disebabkan oleh spirochetes, yaitu Treponema pallidum subspecies pertenue. Penyakit ini merupakan jenis infeksi non venereal kronis dan menular terutama pada anak-anak dengan usia di bawah 15 tahun. Kementerian Kesehatan Republik Indonesia mencatat sebanyak 7.400 kasus frambusia baru dalam periode Oktober 2008 - Oktober 2009 di propinsi Nusa Tenggara Timur (NTT). Diagnosis frambusia sangat memerlukan pemeriksaan serologis sehingga diperlukan metode pemeriksaan yang sederhana, cepat, dan akurat. Rapid Plasma Reagin (RPR) merupakan pemeriksaan penunjang serologis akurat, ekonomis, cepat, dan dapat diulang dengan hasil yang sama.

Tujuan : Mengetahui sensitivitas, spesifisitas, nilai prediksi positif (NPP), dan prediksi negatif (NPN) RPR sebagai penunjang serologis untuk diagnosis frambusia dibandingkan dengan TPHA sebagai baku emas diagnostik frambusia pada anak usia 1-5 tahun.

Subyek dan metode : Penelitian ini merupakan uji diagnostik. Subyek penelitian (SP) adalah sebagian dari anak berusia 1 - 5 tahun di kecamatan Kodi dan Kodi Utara, kabupaten Sumba Barat Daya, NTT. Sejumlah 168 SP telah dilakukan anamnesis, pemeriksaan fisis, dan pengambilan spesimen darah. Serum didapatkan melalui proses sentrifugasi pada setiap spesimen yang kemudian disimpan dalam keadaan beku.

Pemeriksaan RPR dilakukan di laboratorium poliklinik Departemen Ilmu Kesehatan Kulit dan Kelamin.

Pemeriksaan TPHA dilakukan di Departemen Patologi Klinik RSUPN Cipto Mangunkusumo Jakarta.

Hasil : Nilai sensitivitas RPR sebesar 77,8%, nilai spesifitas sebesar 94,7%, NPP sebesar 63,6%, NPN sebesar 97,3%, dan nilai akurasi 92,9%. Lokasi lesi yang paling sering didapatkan adalah di tungkai bawah 85,71%. Jenis lesi kulit yang paling sering didapatkan adalah ulkus 42,85%.

Kesimpulan : Dengan hasil-hasil tersebut dapat disimpulkan bahwa pemeriksaan RPR pada anak usia 1 - 5 tahun sebagai pemeriksaan penunjang serologis dalam menegakkan diagnosis frambusia. Jenis dan lokasi lesi tersering yang ditemukan adalah ulkus dan tungkai bawah.

<hr><i>Background : Yaws is an infection caused by spirochetes, which is Treponema pallidum subspecies pertenue. Yaws is an infectious and chronic non-venereal disease, affecting mostly children between one

and five years old. The Indonesian Ministry of Health reported 7,400 new cases of yaws in Nusa Tenggara Province (NTT) between October 2008 and October 2009. Diagnostic of yaws requires serological diagnostic tools. Hence, a simple, accurate and fast was needed. Rapid Plasma Reagin (RPR) was used as a serological diagnostic tool because RPR is considered to be an accurate, fast, cheap, and reliable tool.

Objective : to measure sensitivity, specificity, Positive Prediction Value (PPV), and Negative Prediction Value (NPV) of RPR as a serological diagnostic tool for yaws in children between one and five years old.

Subjects and method : randomized, diagnostic study was conducted among children between one and five years old in Kodi and Kodi Utara sub-districts of Sumba Barat Daya district , NTT province. Anamnesis, physical examination, and blood samples were collected from 168 subjects. Serum was obtained via the centrifugation of each blood sample, after which it was stored in below zero temperature. RPR test was conducted in an outpatient laboratory at the Department of Dermato-venereology while TPHA test was done at the Department of Clinical Pathology at dr. Cipto Mangunkusumo general hospital.

Result : RPR sensitivity result is 77,8%, specificity result is 94,7%, PPV is 63,6%, NPV is 97,3%, accuracy is 92,9%. Lower extremities are the most affected site in 85,71% subjects. Ulcers (42,85%) are the most common skin lesion recorded in this study.

Conclusion : Based from this results, RPR test is a useful serological diagnostic tool for yaws in children between one and five years old. Lower extremities are the most affected site with ulcers as the most common skin lesion recorded.