

Analisis evolusi polusi udara di Jakarta Indonesia berdasarkan metode biplot = Analysis on air pollution evolution in Jakarta Indonesia based on biplot method / Denny Fitrial

Denny Fitrial, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20390010&lokasi=lokal>

Abstrak

ABSTRAK

Data konsentrasi rata-rata bulanan NO₂, SO₂ dan TSP dari tahun 2009 sampai 2012 di tiga stasiun pemantau polusi udara di Jakarta telah dianalisis dengan menggunakan metode Biplot. Tujuannya adalah untuk menentukan hubungan antara polutan, pengaruh musim terhadap polusi udara dan untuk menggambarkan evolusi polusi udara di kota ini. Metode Biplot dipilih untuk menganalisis karena metode ini dapat digunakan untuk menampilkan variabel dan kasus secara bersamaan. Dengan menggunakan metode Biplot, hubungan antar variabel, kesamaan antar kasus dan posisi relatif variabel dan kasus dapat ditunjukkan.

Evolusi polusi udara di Jakarta menunjukkan suatu perubahan kecenderungan dalam tingkat polusi selama periode pengamatan, dimana konsentrasi TSP dan NO₂ cenderung untuk menurun, sementara itu konsentrasi SO₂ cenderung meningkat. Biplot juga menunjukkan bahwa musim tidak mempengaruhi tingkat polusi di Jakarta dan tidak ada korelasi yang signifikan diantara jenis-jenis polutan yang berbeda.

<hr>

ABSTRACT

The data of monthly average concentration of NO₂, SO₂ and TSP from 2009 to 2012 in three air pollution monitoring stations in Jakarta have been analyzed by using Biplot methods. The objective is to determine the relationship among pollutants, the effect of seasons on air pollution and to describe the evolution of air pollution in the city. Biplot method was chosen to conduct the analysis because it could be used to present the variables and cases simultaneously. By using Biplot method, the relationship among the variables, the similarity among the cases and the relative positions of variables and cases can be shown. Evolution of air pollution in Jakarta shows a change of trend in the pollution levels during the observation periods, where the concentration of TSP and NO₂ tends to decrease, while the concentration of SO₂ tends to increase. Biplot also shows that seasons do not affect the air pollution level in Jakarta and there was no significant correlation between the different types of pollutants.