

Analisis hubungan antara penggunaan root surface conditioning terhadap tingkatan penghilangan smear layer evaluasi (Evaluasi Scanning Electron Microscope) = Analysis the relationship between the use of root surface conditioning with the removal of smear layer study of a (Evaluasi Scanning Electron Microscope

Desy Fidyawati, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20403954&lokasi=lokal>

Abstrak

[ABSTRAK

Latar Belakang: Peran root surface conditioning terhadap keberadaan smear layer.

Tujuan: Menganalisis hubungan antara penggunaan root surface conditioning (minosiklin 2,1% dan EDTA 24%) terhadap keberadaan smear layer setelah penghalusan akar gigi.

Metoda: Sepuluh gigi manusia yang dicabut akibat kelainan periodontal dan dilakukan penghalusan akar.

Gigi dipotong pada daerah

sepertiga servikal, dan 30 spesimen yang terbentuk dibagi dalam tiga kelompok.

Hasil: Tidak ada perbedaan signifikan terhadap tingkat keberadaan smear layer antara kelompok minosiklin maupun EDTA ($p=0,759$). Terdapat perbedaan signifikan antara kelompok minosiklin dan EDTA dengan salin sebagai kontrol ($p=0,00$).

Kesimpulan: Ada hubungan antara penggunaan root surface conditioning terhadap keberadaan smear layer.

<hr>

ABSTRACT

Background: Role of root surface conditioning of the existence of smear layer.

Objective: To analyze the smear layer on root surface conditioned with minocycline HCl 2,1% , EDTA gel 24% , after root planed.

Materials and methods: Ten human teeth removed due to chronic periodontitis were collected and root planed. The teeth were sectioned on 1/3 cervical, 30 specimens were divided into three groups. Results: No significant differences of smear layer between minocycline and EDTA ($p=0,759$). There was significant differences of minocycline and EDTA group compare to saline ($p=0,00$).

Conclusion: There was relationship of root surface conditioning treatment with smear layer.;Background:

Role of root surface conditioning of the existence of smear layer.

Objective: To analyze the smear layer on root surface conditioned with minocycline HCl 2,1% , EDTA gel 24% , after root planed.

Materials and methods: Ten human teeth removed due to chronic periodontitis were collected and root planed. The teeth were sectioned on 1/3 cervical, 30 specimens were divided into three groups. Results: No significant differences of smear layer between minocycline and EDTA ($p=0,759$). There was significant differences of minocycline and EDTA group compare to saline ($p=0,00$).

Conclusion: There was relationship of root surface conditioning treatment with smear layer., Background:

Role of root surface conditioning of the existence of smear layer.

Objective: To analyze the smear layer on root surface conditioned with minocycline HCl 2,1% , EDTA gel 24% , after root planed.

Materials and methods: Ten human teeth removed due to chronic periodontitis were collected and root planed. The teeth were sectioned on 1/3 cervical, 30 specimens were divided into three groups. Results: No significant differences of smear layer between minocycline and EDTA ($p=0,759$). There was significant differences of minocycline and EDTA group compare to saline ($p=0,00$).

Conclusion: There was relationship of root surface conditioning treatment with smear layer.]