

Uji Jalan 6 Menit pada Pasien Pasca TB Paru dan Hubungannya dengan Gejala Klinis TB serta Gambaran Foto X-Ray Toraks di Kabupaten Timor Tengah Selatan Nusa Tenggara Timur = Six Minutes Walking Test of Post Pulmonary TB Patients and It's Associated with Clinical Symptoms of TB and Chest X-Ray Findings in South Central Timor District East Nusa Tenggara

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20411703&lokasi=lokal>

Abstrak

Prevalensi TB paru di Indonesia yang tinggi yaitu sebesar 272 per 100.000 penduduk dan dampaknya terhadap sosio-ekonomi serta kualitas hidup penderitanya membuat penyakit ini sebagai salah satu masalah kesehatan nasional. Pasien TB paru mengalami perubahan fungsi paru akibat inflamasi kronik sehingga terjadi penurunan kualitas hidup. Uji jalan 6 menit merupakan salah satu tes sederhana yang telah terstandarisasi untuk menilai kapasitas fungsional paru.

Penelitian ini bertujuan untuk mengetahui hasil uji jalan 6 menit pada pasien pasca TB paru dan hubungannya dengan gejala klinis TB serta gambaran foto X-Ray toraks. Penelitian ini merupakan penelitian cross sectional yang dilakukan di Kabupaten Timor Tengah Selatan, Nusa Tenggara Timur pada Juni 2011. Sampel penelitian berjumlah 78 orang yang dipilih dengan metode total sampling. Dilakukan wawancara untuk mengisi kuesioner, pengukuran uji jalan 6 menit, dan pemeriksaan foto X-Ray toraks. Rerata hasil uji jalan 6 menit pada laki-laki adalah $438,19 \pm 117,77$ m dan pada perempuan adalah $369,56 \pm 143,10$ m, serta hanya 9 orang subyek (11,5%) yang mencapai hasil uji jalan 6 menit yang normal. Sebesar 56,41% subyek masih memiliki gejala klinis TB dan 88,5% memiliki lesi pada gambaran foto X-Ray toraks. Dapat disimpulkan bahwa terdapat hubungan bermakna antara hasil uji jalan 6 menit dengan gejala klinis TB ($p=0,009$) dan gambaran foto X-Ray toraks ($p=0,000$).

.....

The prevalence of pulmonary TB in Indonesia is high (272 in 100.000 populations). It affects socio-economy and quality of life of the patients so TB is one of national health problems. Lung function in post pulmonary TB will decline as a result of chronic inflammation leading to decreased quality of life. The six minutes walking test is one of standardized simple tests to assess the functional capacity of the lungs. This study aims to determine the result of six minutes walking test of the post pulmonary tuberculosis patients and whether it is associated with clinical symptoms and chest X-Ray findings of TB. This is a cross sectional study held in South Central Timor District, East Nusa Tenggara on June 2011. Seventy eight subjects were selected using total sampling and interviewed to find out any clinical symptoms left. Then, the patients were ask to complete six minutes walking test measurement and chest X-Ray examination. The mean result of six minutes walking test for male is 438.19 ± 117.77 m and for the female is 369.56 ± 143.10 m. Nine out of 78 subjects (11.5%) achieve normal results. Percentage of subjects who still have clinical symptoms of TB is 56.41% and 88.5% shows lesions on chest X-Ray. It is concluded that there is a correlation between six minutes walking test result with the clinical symptoms of TB ($p = 0.009$) and chest X-Ray findings ($p = 0.000$).