

Faktor dominan terhadap kejadian kegemukan pada siswa sma islam al azhar 2 pejaten jakarta selatan tahun 2015 = Dominant factor of overweight among student at al azhar 2 islamic high school pejaten south jakarta 2015

Annisa Isnaeni, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20412088&lokasi=lokal>

Abstrak

Kegemukan merupakan masalah kesehatan yang sering terjadi pada remaja. Penelitian ini bertujuan untuk mengetahui faktor dominan terhadap kejadian kegemukan pada siswa SMA Islam Al-Azhar 2 Pejaten Jakarta Selatan tahun 2015. Penelitian ini menggunakan desain studi cross sectional dengan jumlah sampel 128 siswa. Penelitian ini dilakukan pada bulan April hingga Mei 2015. Pengumpulan data dilakukan melalui pengukuran antropometri (meliputi berat dan tinggi badan), kuesioner mandiri dan wawancara food recall 3x24 jam. Adapun variabel yang diteliti adalah indeks massa tubuh menurut umur (IMT/U), jenis kelamin, asupan zat gizi, aktivitas fisik, kebiasaan konsumsi makanan cepat saji, kebiasaan konsumsi minuman manis, kebiasaan sarapan serta pengetahuan gizi.

Hasil penelitian menunjukkan prevalensi kegemukan sebesar 31,2% dan terdapat hubungan yang signifikan antara jenis kelamin, asupan serat dan kebiasaan konsumsi minuman manis dengan kejadian kegemukan (p -value < 0,05), dimana kebiasaan konsumsi minuman manis merupakan faktor dominan kegemukan.

<hr><i>Overweight is a common health problem in adolescent. This study aimed to identify the dominant factor associated with overweight among student at Al-Azhar 2 Islamic High School Pejaten, South Jakarta 2015. This study used a cross-sectional design with total sample 128 students. This research was conducted in April until May 2015. The data were collected by anthropometry measurements (consist of weight and height), self-administered questionnaire and food recall 3x24 hours interview. The variables studied were BAZ, sex, nutrient intake (energy, protein, carbohydrate, fat and fiber intake), physical activity, fast food consumption, sweetened beverages consumption, breakfast habit and nutrition knowledge.

The result showed that the prevalence of overweight was 31,2%, and there was a significant correlation between sex, fiber intake and consumption of sweetened beverages with overweight (p -value < 0,05) and consumption of sweetened beverages as the dominant factor of overweight.</i>