

Hubungan karakteristik individu dan pelayanan kb dengan penggunaan kontrasepsi suntikan di Indonesia analisis sdki 2012 = Relationship between individual characteristics and family planning services with the use of contraceptive injection in Indonesia 2012 secondary data analysis of 2012 sdki

Kanya Anindya, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20412279&lokasi=lokal>

Abstrak

Kontrasepsi suntikan merupakan alat kontrasepsi yang paling banyak digunakan di tahun 2012. Meski memiliki beberapa keuntungan, kontrasepsi suntikan memiliki berbagai efek samping bagi kesehatan akseptor KB. Penelitian dilakukan untuk menganalisis hubungan karakteristik individu dan pelayanan KB dengan penggunaan kontrasepsi suntikan pada akseptor KB di Indonesia tahun 2012. Penelitian menggunakan desain studi cross sectional dan data Survei Demografi Kesehatan Indonesia Tahun 2012. Hasil analisis multivariabel menunjukkan bahwa seluruh variabel dari faktor karakteristik individu dan pelayanan KB memiliki hubungan yang signifikan dengan penggunaan kontrasepsi suntikan. Diketahui pula bahwa biaya kontrasepsi yang tinggi di pelayanan swasta adalah faktor yang paling dominan yang mempengaruhi penggunaan kontrasepsi suntikan (OR=9,7; nilai p=0,000).

.....Contraceptive injection is the most popular contraception method in 2012. This method offers several advantages, but on the other hand, it also has many side effects for acceptors's health. The study purpose is to analyze the relationship between individual characteristics and family planning services with the use of injectable contraceptives in family planning acceptors in Indonesia in 2012. This study uses cross sectional study design and secondary data analysis of 2012 Indonesia Demographic Health Survey.

The results of multivariable analysis showed that both individual characteristics and family planning services factors have a significant association with the use of contraceptive injection. It also found that the high cost of contraceptives in private services is the most dominant factor that affecting the use of contraceptive injections (OR=9,7; p-value=0,000).