

Perbaikan pola sirkadian tekanan darah donor ginjal dengan ambulatory blood pressure monitoring pada 12 minggu pasca nefrektomi = Normalization of blood pressure circadian rhythm in kidney donors within 12 weeks post nephrectomy

Hasan Basri, 1967-, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20417131&lokasi=lokal>

Abstrak

[**ABSTRAK**] Latar Belakang : Isu yang berkembang pada donor ginjal hidup adalah penurunan fungsi ginjal dan terjadinya hipertensi setelah dilakukan nefrektomi. Satu minggu setelah nefrektomi pola tekanan darah sirkadian berubah menjadi non dipper. Selanjutnya terjadi kompensasi sehingga fungsi ginjal akan stabil dalam 12 minggu. Namun belum diketahui apakah perbaikan fungsi ginjal akan diikuti oleh pola tekanan darah sirkadian kembali menjadi dipper.

Tujuan : Untuk mengetahui perubahan pola sirkadian tekanan darah donor ginjal hidup setelah 12 minggu nefrektomi unilateral.

Metode Penelitian : Studi Pre-experimental dengan before and after design. Subyek sebanyak 18 orang donor ginjal hidup sehat yang berusia 18-50 tahun . Peneltian dilakukan di RSCM pada bulan Januari 2015 sampai dengan Mei 2015. Tekanan darah diukur dengan 24 jam ABPM . Pemeriksaan kreatinin darah, eLFG epi dan uACR dilakukan sebelum nefrektomi, pada 1 minggu dan 12 minggu setelah nefrektomi.

Hasil :Terdapat 18 subyek yang memiliki pola dipper sebelum dilakukan nefrektomi unilateral. Tujuh belas subyek mengalami pola non dipper setelah 1 minggu nefrektomi. Pada 12 minggu setelah nefrektomi 16 diantaranya kembali menjadi pola dipper yang bermakna secara statistik ($p<0.001$).

Simpulan : Terdapat perubahan pola sirkadian tekanan darah non dipper kembali menjadi pola dipper pada donor ginjal hidup 12 minggu setelah nefrektomi unilateral.
ABSTRACT
Background : The issue of post nephrectomy in living kidney donor is kidney function decrease and hypertension. One week after nephrectomy circadian pattern of blood pressure becomes non dipper. Then there will be a compensatory of renal function that becomes stable within 12 weeks after nephrectomy. However, whether the improvement of renal function is followed by the circadian pattern of blood pressure becomes dipper is still unknown.

Aims : To know the changes circadian pattern of blood pressure among living kidney donors 12 weeks after unilateral nephrectomy.

Methods : A pre-experimental study with before and after design. The subjects were 18 healthy living kidney donors aged 18 to 50 years old , conducted in RSCM hospital between January 2015 to May 2015. Blood pressure was measured by 24 hours ABPM. Serum creatinine, e-GFR epi and uACR were taken before nephrectomy, 1 week and 12 weeks after nephrectomy.

Results : There were 18 subjects had dipper pattern before unilateral nephrectomy. Seventeen of them exhibited a pattern became non dipper on one week after nephrectomy. Sixteen subjects showed the pattern returned to dipper after 12 weeks nephrectomy that statistically significant ($p<0.01$)

Conclusions : The circadian pattern of blood pressure returned to dipper from non dipper on living kidney donors after 12 weeks unilateral nephrectomy., Background : The issue of post nephrectomy in living kidney donor is kidney function decrease and hypertension. One week after nephrectomy circadian pattern of blood pressure becomes non dipper. Then there will be a compensatory of renal function that becomes stable within 12 weeks after nephrectomy. However, whether the improvement of renal function is followed by the circadian pattern of blood pressure becomes dipper is still unknown.

Aims : To know the changes circadian pattern of blood pressure among living kidney donors 12 weeks after unilateral nephrectomy.

Methods : A pre-experimental study with before and after design. The subjects were 18 healthy living kidney donors aged 18 to 50 years old , conducted in RSCM hospital between January 2015 to May 2015. Blood pressure was measured by 24 hours ABPM. Serum creatinine, e-GFR epi and uACR were taken before nephrectomy, 1 week and 12 weeks after nephrectomy.

Results : There were 18 subjects had dipper pattern before unilateral nephrectomy. Seventeen of them exhibited a pattern became non dipper on one week after nephrectomy. Sixteen subjects showed the pattern returned to dipper after 12 weeks nephrectomy that statistically significant ($p<0.01$)

Conclusions : The circadian pattern of blood pressure returned to dipper from non dipper on living kidney donors after 12 weeks unilateral nephrectomy.]