

Hubungan antara infeksi Soil Transmitted Helminths dengan kejadian Anemia pada balita di Kecamatan nangapanda Ende Nusa Tenggara Timur = Correlation between Soil Transmitted Helminths infection and the prevalence of anemia in under five children in Nangapanda Ende District East Nusa Tenggara

Sitohang, Agrianti, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20421189&lokasi=lokal>

Abstrak

Anemia adalah suatu kondisi dimana jumlah sel darah merah atau kemampuannya untuk mengangkut oksigen tidak cukup untuk memenuhi kebutuhan fisiologis tubuh. Anemia pada anak dapat mempengaruhi tumbuh kembang anak, serta menyebabkan anak menjadi lemas dan pucat. Salah satu faktor yang dapat menyebabkan anemia di negara-negara berkembang adalah infeksi soil-transmitted helminths. Penelitian ini bertujuan untuk meneliti hubungan antara infeksi soil-transmitted helminths dengan kejadian anemia pada balita di Kecamatan Nangapanda, Ende, Nusa Tenggara Timur. Penelitian ini menggunakan desain potong-lintang yang meneliti status infeksi soil-transmitted helminths melalui pemeriksaan telur pada feses dan hubungannya dengan anemia yang diukur dari kadar hemoglobin darah kapiler. Selain itu juga diteliti hubungan antara anemia dengan faktor perancu berupa usia, jenis kelamin, dan nutrisi. Penelitian ini dilakukan pada Juli 2015 hingga Oktober 2015 di Departemen Parasitologi Fakultas Kedokteran Universitas Indonesia. Data dianalisis dengan menggunakan program SPSS 20.0 for windows dengan Uji Chi Square dan Fisher untuk analisis univariat serta Regresi Logistik untuk analisis multivariat. Hasil penelitian ini menunjukkan prevalensi infeksi soil-transmitted helminths pada 100 balita di Kecamatan Nangapanda, Ende, Nusa Tenggara Timur adalah 59%, dengan infeksi terbanyak disebabkan oleh infeksi majemuk antara *Ascaris lumbricoides* dan *Trichuris trichiura* sebanyak 23%. Prevalensi anemia pada balita di Kecamatan Nangapanda, Ende, Nusa Tenggara Timur adalah 63%. Tidak terdapat hubungan yang signifikan antara infeksi soil-transmitted helminths dengan kejadian anemia pada balita di Kecamatan Nangapanda, Ende, Nusa Tenggara Timur. Dari berbagai faktor yang diteliti, usia yang kurang dari 2 tahun merupakan faktor yang memiliki hubungan yang signifikan terhadap kejadian anemia pada populasi ini.

Anemia is a condition where the red blood cell count or its ability to carry oxygen is not adequate to fulfill physiological needs. Anemia in children can affected growth and development and cause fatigue, paleness. One of the factors that can cause anemia in developing countries is soil-transmitted helminths infection. This research aimed to analyze the correlation between soil-transmitted helminths infection and the prevalence of anemia in under-five children in Nangapanda Subdistrict, Ende district, East Nusa Tenggara. This research used cross-sectional design by investigating soil-transmitted helminths infection status through detection of helminth eggs in stool examination and its correlation with status of anemia based on hemoglobin measured from finger-prick blood. The link between anemia and confounding factors such as age, gender, and nutriotional status was also analyzed. This research was conducted from July 2015 to October 2015 at Parasitology Departement of Faculty of Medicine of University of Indonesia. The data are analyzed using SPSS 20.0 software for Windows with Chi-Square and Fisher Test for univariate analysis and Logistic Regression for multivariate analysis. The results showed that the prevalence of soil-transmitted helminths infection in the 100 under-five children in Nangapanda Subdistrict, Ende district, East Nusa

Tenggara was 59%, with majority of children (23%) co infected with *Ascaris lumbricoides* and *Trichuris trichiura*. The prevalence of anemia in the under-five children in Nangapanda Subdistrict, Ende district, East Nusa Tenggara 63%. There was no significant correlation between soil-transmitted helminths infection and anemia status in under-five children in Nangapanda Subdistrict, Ende District, East Nusa Tenggara. However, among other confounding factors analyzed, children under 2 years old were significantly associated with the prevalence of anemia in this population.