

# Distribusi polimorfisme gen wingless type 3a (Wnt3a) rs 752107 pada penderita orofacial cleft = Distribution of wingless type 3a (Wnt3a) rs 752107 gene polymorphism in orofacial cleft

Simbolon, Uji Tesli Haralini Br., author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20421512&lokasi=lokal>

---

## Abstrak

Latar Belakang: Gen Wnt3a berperan pada pembentukan orofacial cleft.

Tujuan: Melihat distribusi polimorfisme gen Wnt3a rs 752107 pada penderita OFC.

Metode: Menggunakan teknik PCR-RFLP pada 30 sampel DNA penderita OFC dan 70 DNA kontrol, diperoleh gambaran polimorfisme gen tersebut.

Hasil: 100% sampel penderita OFC memiliki genotip CC (wildtype). Selanjutnya, dengan uji chi-square menunjukkan tidak terdapat perbedaan bermakna antara distribusi polimorfisme gen Wnt3a rs 752107 pada penderita OFC dan kontrol ( $p>0.05$ ).

Kesimpulan: Pada penelitian ini tidak ditemukan distribusi polimorfisme gen Wnt3a rs 752107 pada penderita OFC.

<hr><i>Background: Wnt3a plays role in the process orofacial cleft.

Aim: to observe the distribution of Wnt3a rs 752107 gene polymorphism in OFC.

Methods: This research were used PCR-RFLP technique of 30 OFC and 70 control DNA samples, obtained a description of the gene polymorphism.

Results: 100% OFC samples have CC genotype (wildtype). Then, with chi-square test in SPSS 22 there were found no significant differences between the distribution of the gene polymorphism in OFC and control ( $p>0.05$ ).

Conclusion: In this research found no distribution of Wnt3a rs 752107 gene polymorphism in orofacial cleft.</i>