

Asuhan keperawatan kesehatan masyarakat perkotaan dengan latihan kegel pada pasien kanker rektum post operasi ultra low anterior resection = Urban health nursing care with kegel exercises on rectum cancer patients post operation ultra low anterior resection / Ihda Fakhriyana Istikarini

Ihda Fakhriyana Istikarini, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20434730&lokasi=lokal>

Abstrak

ABSTRAK

Kanker rektum merupakan salah satu penyakit keganasan saluran gastrointestinal yang banyak dialami masyarakat perkotaan. Salah satu tindakan pembedahan yang dapat dilakukan adalah operasi metode ultra low anterior resection. Berdasarkan jurnal terkait, pasien paksa operasi metode ultra low anterior resection memiliki risiko sebesar 10-20% terkena sindrom anterior resection dengan gejala kelemahan sfingter anal sehingga mengakibatkan inkontinensia fekal. Salah satu tindakan yang dapat dilakukan kepada pasien untuk mencegah masalah tersebut adalah dengan latihan kegel. Latihan kegel yang dilakukan secara rutin dapat membantu meningkatkan kontrol anus dan menguatkan sfingter anus. Latihan dilakukan secara bertahap sebanyak 4x10 set dalam sehari. Evaluasi latihan kegel dilakukan menggunakan colok dubur untuk mengevaluasi kekuatan kontraksi anus. Sebelum latihan kegel dilakukan, penanganan manajemen nyeri pada pasien paska operasi harus dilaksanakan dengan baik terlebih dahulu agar toleransi latihan kegel tinggi dan latihan dapat dilakukan segera setelah operasi.

<hr>

ABSTRAK

Rectal cancer is a malignancy of the gastrointestinal tract that is experienced by the urban community. One of the surgery choices is ultra low anterior resection method. Based on the relevant journal, patients were treated with this method have 10-20% higher risk of anterior resection syndrome, which symptoms is weakness of the anal sphincter, resulting in fecal incontinence. One of the interventions that can prevent patients from the fecal incontinence is Kegel exercises. Regular exercises can help improve and strengthen control of the anal sphincter. Exercises done gradually as 4x10 sets in a day. The evaluation of exercises using a rectal tusche to evaluate the strength of anus contraction. Before Kegel exercises is started, handling the management of pain post-surgery must be performed well so that patients have a higher tolerance of Kegel exercises and the exercises can be done soon after surgery.;