

Optimalisasi pemenuhan kebutuhan cairan dan elektrolit anak melalui deteksi dini plebitis menggunakan skala infusion nurse society dengan pendekatan model konservasi levine = Optimizing of children's fluid and electrolyte fullfillment through early detection using infusion nurse society phlebitis scale with levine s conservation model approach

Diah Ayu Agustin, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20435244&lokasi=lokal>

Abstrak

Hampir 100% anak yang dirawat di rumah sakit terpasang infus untuk mengatasi masalah ketidakseimbangan cairan dan elektrolit. Cairan dan elektrolit merupakan bagian konservasi energi yang dibutuhkan anak untuk mempertahankan fungsi tubuh secara utuh. Model Konservasi Levine berfokus pada peningkatan adaptasi melalui prinsip konservasi untuk mencapai keutuhan diri. Karya ilmiah ini membahas proses keperawatan terhadap lima orang anak yang terpasang infus. Salah satu tindakan konservasi energi untuk optimalisasi pemenuhan kebutuhan cairan dan elektrolit anak adalah deteksi dini plebitis menggunakan skala Infusion Nurse Society dengan pendekatan model konservasi Levine. Hasil evaluasi kelima kasus yang mengalami masalah cairan dan elektrolit menunjukkan 4 kasus masalah teratasi dan satu kasus belum teratasi karena klien mengalami perburukan saat hendak pulang. Deteksi dini plebitis turut berperan penting mempertahankan kelancaran akses intravena untuk terapi cairan dan elektrolit.

Almost 100% of children admitted who inserted infusion to overcome the problem of fluid and electrolyte imbalance. Fluid and electrolyte are part of children energy conservation need to adapt to maintain the body functions as a whole. Levine's conservation model focuses on improving adaptation through conservation principles to achieve wholeness. One of energy conservation intervention to optimize of Children's Fluid and Electrolyte Fulfillment through Early Detection using Infusion Nurse Society Phlebitis Scale with Levine's Conservation Model Approach. This paper discusses the nursing process to the five children who inserted infusion. The results of the evaluation of the five cases that have problems of fluid and electrolyte showed 4 cases the problem is resolved and the case is not resolved because the client experience worsening when going home. Early detection of phlebitis has an important role in maintaining intravenous access for fluid and electrolyte therapy.