

## Penatalaksanaan kasus gangguan sendi temporomandibula pada pasien dengan dimensi vertikal yang rendah dan gigitan anterior yang dalam

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20436057&lokasi=lokal>

---

### Abstrak

Indonesian Journal of Dentistry 2006; Edisi Khusus KPPIKG XIV: 155-158

The causes of TMD are complex and multifactorial, therefore the management should be done by several disciplines. In this report, a 27-year-old man came to the teaching hospital of the University of Indonesia Faculty of Dentistry's Prosthodontic Department complaining about clicking sound and pain around his right joint. He received orthodontic treatment 9 years ago with removable appliance at a private practice and had 4 premolar extraction. The patient's face looked asymmetric, with a low vertical dimension, a Class II occlusion, and an anterior deep bite. Besides that, he clenched his teeth during emotional stress. Lateral tanscranial photo showed that the position of the left condyle was relatively normal or slightly anterior, and the right condyle was in the superoposterior position in the fossa with an abnormal shape. To solve this problem, the patient was referred to the Orthodontic Department to get a correct vertical dimension and normal anterior overbite. After 6 years, the patient was again referred to the prosthodontic Department, but the result was not successful. In order to get the right vertical dimension, an occlusal splint was fabricated to achieve a comfort jaw relation. In this position, the overbite was 2 mm, but space between the upper and lower posterior teeth was 5 mm. In this situation, full veneer crowns were not impossible to fabricate. Finally, to maintain this comfort position, the patient was suggested to wear the occlusal splint and come regularly for control every 6 months.