

**Kejadian ototoksitas pada kasus keganasan anak yang mendapatkan terapi carboplatin dengan pemeriksaan otoacoustic emission (OAE) =  
The incidence of ototoxicity in child malignancy cases who received carboplatin therapy with otoacoustic emission (OAE) examination /  
Jarot Kunto Wibowo**

Jarot Kunto Wibowo, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20445678&lokasi=lokal>

---

**Abstrak**

**<b>ABSTRAK</b><br>**

Latar belakang: Keganasan merupakan salah satu penyakit tidak menular yang menjadi masalah kesehatan masyarakat, baik di dunia maupun di Indonesia. Salah satu modalitas pengobatan kasus keganasan adalah kemoterapi. Carboplatin cis-diammine-cyclobutanedicarboxylato platinum adalah senyawa platinum generasi kedua yang sering digunakan dalam tata laksana kasus keganasan seperti neuroblastoma, retinoblastoma, hepatoblastoma, tumor otak dan tumor sel germ. Efek samping pemberian obat sitotoksik perlu dipertimbangkan, khususnya ototoksik, yaitu gangguan fungsi dan kerusakan struktur telinga dalam yang dapat disebabkan oleh obat atau bahan kimia tertentu. Tujuan: Menilai efek ototoksitas akibat pengaruh Carboplatin pada anak dengan kasus keganasan. Metode: Penelitian ini menggunakan desain serial cross sectional untuk mengetahui perubahan signal to noise ratio SNR pada OAE sebagai akibat efek ototoksik dan faktor-faktor risiko yang ikut berperan pada kejadian ototoksik akibat pemakaian Carboplatin di Divisi Hematologi Onkologi IKA FKUI-RSCM. Hasil: terdapat dua dari 52 subyek penelitian yang mengalami kejadian ototoksik. Kesimpulan: didapatkan dua 5 dari 40 subyek mengalami kejadian ototoksik pada kelompok yang mendapat kemoterapi, sedangkan pada kelompok yang belum mendapat terapi tidak ditemukan adanya nilai SNR kurang dari enam. Faktor risiko berupa jenis kelamin, usia, siklus kemoterapi dan dosis Carboplatin tidak memiliki hubungan yang bermakna secara statistik  $p>0,05$ . Kata kunci: Carboplatin, ototoksik, otoacoustic emission OAE <hr />

**<b>ABSTRACT</b><br>**

Background Malignancy is one of the non contagious diseases are a public health problem, both globally and Indonesia. Chemotherapy is one of modality in malignancy cases. Carboplatin cis platinum diammine cyclobutanedicarboxylato is a second generation platinum compound that is often used in the management of cases of malignancies such as neuroblastoma, retinoblastoma, hepatoblastoma, brain tumors and germ cell tumors. Side effects of cytotoxic drugs need to be considered, especially ototoxic. Ototoxic is dysfunction and damage to the structure of the inner ear that can be caused by drugs or other certain chemicals. Objective Assess the effects of ototoxicity due to the influence of Carboplatin in children with malignancy cases. Methods This study uses a serial cross sectional design to assess changes in signal to noise ratio SNR at the OAE as a result of ototoxic effects and risk factors that come into play in the event due to the use of ototoxic Carboplatin in the Hematology Oncology of pediatric Department Cipto Mangunkusumo General Hospital. Results There are two of 52 study subjects, who experienced ototoxic. Conclusion we obtained two 5 of the 40 subjects experienced ototoxic events in the group receiving chemotherapy, whereas in the group that had not received therapy was not found SNR value less than six. The risk factors such as gender, age, Carboplatin dose cycles of chemotherapy and did not have a

statistically significant relationship p 0.05 .