

Hubungan riwayat merokok dan keberhasilan pengobatan fase intensif penderita tuberculosis paru di RSUD dr Zainoel Abidin Banda Aceh = The Association between smoking history and intensive phase treatment success of pulmonary tuberculosis patients in dr Zainoel Abidin hospital Banda Aceh

Risa Fitria, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20447451&lokasi=lokal>

Abstrak

ABSTRAK

Latar belakang : Banyak penyakit yang dihubungkan dengan merokok seperti penyakit keganasan, kardiovaskuler, diabetes mellitus DM , penyakit paru obstruktif kronik PPOK , arthritis, impotensi, infertilitas, alzheimer, TB dan lain- lain. Paru merupakan organ yang banyak mengalami kerusakan berat akibat merokok. Merokok terbukti mengganggu bersihan mukosilier dan berhubungan dengan peningkatan risiko terjadinya TB paru. Konversi sputum merupakan indikator penting untuk melihat keberhasilan pengobatan TB. Penelitian ini bertujuan untuk mengetahui bagaimana hubungan riwayat merokok dan keberhasilan pengobatan fase intensif tuberculosis TB paru di RSUD. Metode : Desain penelitian menggunakan metode kohort prospektif yang dilakukan pada pasien TB paru basil tahan asam BTA positif perokok dan bukan perokok yang berkunjung ke Pelayanan Tuberculosis Terpadu PTT yaitu poli rawat jalan dan ruang rawat inap infeksi paru RSUD Dr. Zainoel Abidin Banda Aceh dari 1 November 2015 hingga Februari 2016. Data yang diperoleh diuji dengan menggunakan statistical package for social sciences SPSS 20. Hasil : Total 38 subjek dibagi kedalam 2 kelompok 19 subjek perokok dan 19 subjek bukan perokok , semua subjek perokok adalah laki-laki sedangkan subjek bukan perokok terdiri dari laki-laki dan perempuan. Dari hasil penelitian bulan pertama $p=0,009$ didapatkan subjek yang bukan perokok konversi sputum BTA berjumlah 14 orang 73,7 dan 5 orang tidak konversi, sedangkan pada subjek perokok 6 orang 31,6 yang konversi dan 13 orang 68,4 tidak konversi. Pada bulan kedua $p=0,202$, lebih dari setengah jumlah subjek yang bukan perokok konversi sputum BTA berjumlah 17 orang 89,5 dan 2 orang 10,5 yang tidak konversi sedangkan pada subjek perokok yang konversi berjumlah 14 orang 73,7 dan 5 orang 26,3 yang tidak konversi. Kesimpulan : Terdapat hubungan bermakna antara kebiasaan merokok terhadap konversi sputum BTA bulan I. Kata kunci : TB paru, merokok, konversi sputum

<hr>

ABSTRACT

Background Many diseases are associated with smoking such as malignant disease, cardiovascular, diabetes mellitus DM , chronic obstructive pulmonary disease COPD , arthritis, impotence, infertility, Alzheimer Disease, tuberculosis and others. Lung is an organ that suffered heavy damage from smoke. Smoking is proven to disrupt the ciliary mucosal clearance and it is associated with an increased risk of pulmonary tuberculosis. Sputum conversion is an important indicator to assess the success of TB treatment. This study aims to determine the relation between smoking history and the success intensive phase treatment of pulmonary tuberculosis TB at Dr Zainoel Abidin Hospital. Method This is a prospective cohort study in patients with pulmonary tuberculosis acid fast bacilli AFB positive smokers and non smokers who visited the Integrated Tuberculosis Care PTT , at outpatient and inpatient pulmonary infection RSUD Dr. Zainoel

Abidin hospital Banda Aceh from 28 November 2015 until 1 February 2016. The data were tested using statistical package for social sciences SPSS 20. Results A total of 38 subjects were divided into 2 groups 19 subjects with 19 subjects smokers and non smokers . All subjects smokers are male while non smoker subjects consisted of male and female. The results of the first month study $p = 0.009$, there are 14 non smoker subjects with AFB conversion 73,7 and 5 subjects without AFB conversion. Among smoking subjects there are 6 subjects 31.6 with AFB conversion and 13 subjects 68.4 without AFB conversion. In the second month $p = 0,202$, more than half subjects who are non smokers had AFB conversion, 17 subjects 89.5 and 2 subjects 10.5 had no AFB conversion. In smokers group there are 14 subjects 73.7 had AFB conversion and 5 subject 26.3 had no conversion. Conclusion There was a significant relation between smoking habit and the occurrence of first month AFB sputum