

Faktor-faktor yang berhubungan dengan anemia maternal dan penyebabnya di RSUD AM Parikesit Tenggarong tahun 2017 = Factors related to maternal anemia and its causes in RSUD AM Parikesit Tenggarong 2017

Herlena Hayati, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20454590&lokasi=lokal>

Abstrak

Menurut WHO 2008, Indonesia termasuk negara dengan prevalensi anemia katagori berat ge;40 , Riskesdes 2013 menyebutkan prevalensi anemiamaterna sebesar 37,1 . Perdarahan merupakan penyebab utama kematian ibu 30.3 dan anemia maternal merupakan faktor risiko utama perdarahan. 51,5 kematian maternal disebakan oleh anemia maternal sebagai penyebab kematiantidak langsung. Penelitian ini bertujuan untuk mengetahui faktor-faktor yangberhubungan dengan anemia maternal dan penyebabnya di RSUD AM ParikesitTenggarong Tahun 2017. Penelitian menggunakan metode penelitian kuantitatif dan kualitatif. Rancangan kuantitatif menggunakan metode deskriptif analitik dengan desain studi cross sectional dengan jumlah sampel 214 ibu hamil. Sedangkan rancangan kualitatif menggunakan RAP, analisis data menggunakan content analisis dengan jumlah informan 37 orang. Hasil penelitian menunjukkan ada hubungan yang bermakna antara status gizi RP=1,7; 95 CI: 1,1-2,6 dengan anemia maternal setelah di kontrol kepemilikan rumah, riwayat pertolongan persalinan, jarak kehamilan dan paritas. Tidak rumah sendiri RP=1,5; CI: 1,0-2,2 berisiko anemia. Riwayat pertolongan persalinan sebelumnya bidan RP=0,6; 95 CI: 0,4-0,9 berisiko lebih rendah untuk anemia maternal.

Hasil penelitian kualitatif menunjukkan konsumsi heme nabati adalah konsumsi proteinterbanyak pada ibu anemia selain rendahnya kepatuhan ibu hamil dalam mengkonsumsi tablet Fe dan pemahaman yang salah tentang penyebab anemiamaterna pada kelompok ibu hamil. Hal yang sebaliknya ditemukan pada kelompok ibu hamil yang tidak anemia maternal, konsumsi heme hewani adalah konsumsi harian informan, hampir seluruhnya patuh dalam mengkonsumsi tablet Fe dan menyebutkan kurang asupan makanan bergizi sebagai penyebab terbanyak kurang darah pada ibu hamil. Perlu kebijakan skrening anemia pada remaja dan tablet Fe prakehamilan pada calon pengantin yang anemia termasuk penanganan gizi lintas program dan lintas sektoral. Kata Kunci: Anemia Maternal, Status Gizi, Ibu Hamil.

<hr /><i>According to WHO 2008 , Indonesia is a country with a severe category anemia prevalence ge 40 , Riskesdes data 2013 found 37,1 prevalence of maternal anemia. Bleeding is a major cause of maternal death 30.3 and maternal anemia is a major risk factor for bleeding. 51.5 of maternal deaths are caused by maternal anemia as the cause of indirect death. This study aims to determine the factors related to maternal anemia and its causes in RSUD AM ParikesitTenggarong Year 2017. The research used quantitative and qualitative research methods. Quantitative design uses descriptive analytic method with cross-sectional study design with 214 samples of pregnant women. While the qualitative design using RAP, analysis method used content analysis with number of informant 37 people. The results showed that there was a significant correlation between nutritional status RP 1.7, 95 CI 1.1 2.6 with maternal anemia after controlled home ownership, history of delivery assistance, distance of pregnancy and parity. Not having a home RP 1.5 CI 1.0 2.2 is more at risk of anemia. History of delivery assistance midwives RP 0.6, 95 CI 0.4 0.9 lower risk for maternal anemia.

The result of qualitative research shows that vegetable hemeconsumption is the most protein consumption in the mother of anemia besides thelow compliance of pregnant mother in consuming Fe tablet and the wrongunderstanding about maternal anemia cause in pregnant women group. Theopposite is found in the group of pregnant women who are not maternal anemias,animal heme consumption is the daily consumption of informants, almost entirelyobedient in consuming Fe tablets and mention less intake of nutritious foods asthe most cause of less blood in pregnant women. Anemia screening policy isrequired on adolescents and Fe tablets for fre conception in bridal candidatesincluding cross program and cross sectoral nutritional care.Key words Maternal Anemia, Nutrional Status, Pregnant Women.</i>