

Analisis praktik klinik keperawatan kesehatan masalah perkotaan : batuk efektif, positioning, dan inhalasi dalam mengatasi masalah bersihan jalan nafas pada pasien closed pathological fracture bilateral dengan community acquired pneumonia = Clinical practice analysis of the health urban nursing issues effective cough positioning and inhalation to help airway clearance problem in patients who has closed pathological fracture bilateral with community acquired pneumonia

Ayu Tiarno Lestari, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20459783&lokasi=lokal>

Abstrak

**ABSTRAK
**

Pneumonia adalah penyakit infeksi yang menular melalui udara. Karakteristik kota yang memiliki kepadatan penduduk, pemukiman dan transportasi, membuat penyebaran pneumonia menjadi lebih mudah. Salah satu masalah keperawatan pada pneumonia adalah ketidakefektifan bersihan jalan nafas. Studi kasus ini bertujuan untuk menganalisis pemberian intervensi yaitu batuk efektif, positioning, dan inhalasi pada klien dengan Closed Pathological Fracture Bilateral dengan Community Acquired Pneumonia. Hasil yang diperoleh yaitu pengeluaran sekret yang kurang maksimal berhubungan dengan keterbatasan pasien dalam melakukan mobilisasi dan positoning karena nyeri akibat fraktur. Rekomendasi untuk intervensi selanjutnya adalah pemberian posisi semi fowler 45 atau tegak lurus harus diupayakan agar terjadi pengembangan ekspansi paru yang dapat meningkatkan tekanan ekspirasi maksimum yang memudahkan dalam pengeluaran sekret.

<hr />

**ABSTRACT
**

Pneumonia is an airborne infectious disease. Urban is characterized by its density of residents, living area and transportation which make the spreading of infectious disease is easier. One of the nursing problem of Pneumonia is Ineffective Airway Clearance. This case study was conducted to analyze the intervention of the effective cough, positioning and inhalation for client who had closed pathological bilateral fracture with community acquired pneumonia. The results of the intervention showed less optimal in mucous secretion related to client rsquo s immobilization and position because of pain due to fractures. This study recommend nurses should give positioning for 45 semi fowler or sit to maximize the pulmonary expansion which may increase the maximum expiratory pressure which facilitates the discharge of secretions.