

Perbedaan kadar imunoglobulin E spesifik serum kuantitatif akibat sensitisasi dermatophagoides pteronyssinus, dermatophagoides farinae dan blomia tropicalis pada asma alergik intermiten dan persisten = Difference of serum quantitative specific ige level induced by dermatophagoides pteronyssinus dermatophagoides farinae and blomia tropicalis sensitization in intermittent and persistent allergic asthma

Agus Joko Susanto, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20460543&lokasi=lokal>

Abstrak

Latar belakang: Tungau debu rumah TDR merupakan alergen hirup yang penting pada asma alergik. Namun, penelitian diagnostik molekuler menggunakan Imunoglobulin E IgE spesifik akibat sensitisasi alergen TDR dihubungkan dengan derajat keparahan asma alergik belum pernah dilakukan di Indonesia.

Tujuan: Mengetahui perbedaan kadar IgE spesifik serum kuantitatif akibat sensitisasi alergen Dermatophagoides D. pteronyssinus, D. farinae dan Blomia B. tropicalis pada asma alergik intermiten dan persisten.

Metode: Desain penelitian potong lintang pada pasien asma alergik dewasa yang diundang untuk pemeriksaan IgE spesifik serum dan merupakan bagian dari penelitian payung di Divisi Alergi dan Immunologi Klinik, RS Cipto Mangunkusumo. Derajat keparahan asma ditentukan berdasarkan kriteria Global Initiative on Asthma GINA 2015 dan dikelompokkan menjadi intermiten dan persisten. Pemeriksaan IgE spesifik serum kuantitatif menggunakan metode multiple allergosorbent test Polycheck Allergy, Biocheck GmbH, Munster, Germany . Alergen TDR yang diperiksa adalah D. pteronyssinus, D. farinae, dan B. tropicalis. Perbedaan antara dua kelompok dianalisis dengan uji Mann-Whitney.

Hasil: Sebanyak 87 subyek dilibatkan dalam penelitian ini; 69 79,3 subyek adalah perempuan. Rerata usia pasien adalah 40,2 tahun. Enam puluh tiga 72,4 pasien menderita asma dan rinitis alergik. Sebanyak 58 66,7 pasien asma persisten. Gambaran sensitisasi alergen TDR adalah 62,1 D. farinae; 51,7 D. pteronyssinus dan 48,3 B.tropicalis. Median kadar IgE spesifik secara bermakna lebih tinggi pada asma persisten dibandingkan asma intermiten untuk alergen D. farinae 1,30 vs. 0,0 kU/L; $p=0,024$ dan B. tropicalis 0,57 vs. 0,0 kU/L; $p=0,015$. Kadar IgE spesifik D. pteronyssinus lebih tinggi pada asma persisten dibandingkan intermiten 0,67 vs. 0,00 kU/L; $p=0,066$.

Kesimpulan: Gambaran sensitisasi alergen secara berurutan didapatkan D. farinae 62,1, D. pteronyssinus 51,7 dan B. tropicalis 48,3 . Kadar IgE spesifik akibat sensitisasi D. farinae dan B. tropicalis lebih tinggi secara bermakna pada pasien asma persisten dibandingkan asma intermiten. Kadar IgE spesifik akibat sensitisasi D. pteronyssinus lebih tinggi pada pasien asma persisten dibandingkan asma intermiten, tetapi secara statistik tidak bermakna.

<hr>

Introduction House dust mites HDM are an important inhalant allergen in allergic asthma. However,

molecular diagnostic study using specific IgE level induced by HDM allergens associated with asthma severity has not been done in Indonesia.

Objective To investigate the difference of serum quantitative specific IgE levels induced by Dermatophagoides *D. pteronyssinus*, *D. farinae* and *Blomia B. tropicalis* sensitization in intermittent and persistent allergic asthma.

Method This was a cross sectional study on adult allergic asthma patients who were invited for serum specific IgE testing. This study was a part of a larger research within the Division of Allergy and Immunology, Cipto Mangunkusumo Hospital. Asthma severity was defined based on Global Initiative on Asthma GINA 2015 criteria and were grouped as intermittent or persistent. Quantitative specific IgE testing was done on blood serum using a multiple allergosorbent test Polycheck Allergy, Biocheck GmbH, Munster, Germany . The HDM allergens tested were *D. pteronyssinus*, *D. farinae*, and *Blomia tropicalis*. Difference between two groups were analyze using Mann Whitney test.

Results A total of 87 subjects were enrolled in this study 69 79.3 were women. Mean patients rsquo age was 40, 2 years. Sixty three 72.4 patients had asthma and allergic rhinitis. Fifty eight 66.7 patients were classified as persistent asthma. The prevalence of sensitization was 62.1 *D. farinae*, 51.7 *D. pteronyssinus*, and 48.3 *Blomia tropicalis*. The median of specific IgE levels is significantly higher in persistent asthma compares to intermittent asthma induced by *D. farinae* median 1.30 vs. 0.0 kU L p 0.024 and *B. tropicalis* median 0.57 vs. 0.0 kU L p 0.015 sensitization. Level of Specific IgE *D. pteronyssinus* is also to be higher in persistent asthma than the level measured in intermittent asthma 0.67 vs. 0.00 kU L p 0.066.

Conclusion Sensitization of HDM allergens is shown to be highest for *D. farinae* 62.1 , followed by *D. pteronyssinus* 51, 7 and *Blomia tropicalis* 48, 3 . Specific IgE level induced by *D. farinae* and *Blomia tropicalis* sensitization are significantly higher in patients with persistent compares to intermittent asthma, whereas specific IgE level induced by *D. pteronyssinus* sensitization to be higher in persistent asthma although not statistically significant.