

Uji validitas dan reliabilitas University of Washington quality of life questionnaire version 4 adaptasi Bahasa Indonesia penilaian kualitas hidup pasien kanker kepala dan leher pasca terapi 2-6 bulan di departemen THT-KL RSUPN dr. Cipto Mangunkusumo = Validity and reliability test of the Indonesian version of university of Washington quality of life questionnaire version 4 assessment of quality of life of head and neck cancer patient 2-6 months post therapy in ORL HNS Department dr Cipto Mangunkusumo

Karina Anggiaty Idris Gassing, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20460737&lokasi=lokal>

Abstrak

Latar Belakang: Kanker kepala dan leher terdapat 10 dari keseluruhan kasus kanker di seluruh tubuh. Efek samping akibat terapi kanker berdampak signifikan pada kualitas hidup pasien. Instrumen yang sering digunakan untuk menilai kualitas hidup pasien kanker kepala dan leher salah satunya adalah University of Washington Quality of Life UW-QOL. Hingga saat ini belum pernah dilakukan adaptasi kuesioner UW-QOL ke bahasa Indonesia.

Tujuan: Penelitian ini bertujuan mendapatkan instrumen UW-QOL adaptasi bahasa Indonesia yang valid dan reliabel untuk menilai kualitas hidup pasien kanker kepala dan leher.

Metodologi: Penelitian ini berdesain potong lintang, dilakukan di poliklinik THT FKUI/RSCM dr. Cipto Mangunkusumo terhadap pasien kanker kepala dan leher usia dewasa.

Hasil: Uji validitas menggunakan uji korelasi Spearman dengan korelasi bermakna pada seluruh butir pertanyaan di tingkat signifikansi p

.....Background: Head and neck cancer accounts for 10 of all cancer cases throughout the body.. Side effects due to cancer therapy have a significant impact on patient quality of life. The University of Washington Quality of Life UW QOL is the most frequent instruments used to assess the quality of life of head and neck cancer patients. At present, the Indonesian version of UW QOL questionnaire is not available.

Objective: This study aims to obtain a valid and reliable Indonesian adaptation of UW QOL to assess the quality of life of head and neck cancer patients.

Method: Cross sectional study was conducted in ORL HNS Department outpatient clinic dr. Cipto Mangunkusumo hospital towards 41 adult patients with head and neck cancer.

Result: The validity test using Spearman correlation test with significant correlation in all questions items at the level of significance p