

## Shale hydrocarbon potential of talang akar and lemat/lahat formations in south and central Palembang sub-basin / Julikah, Sriwijaya, Jonathan S., Panuju

Julikah, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20471376&lokasi=lokal>

---

### Abstrak

#### **ABSTRACT**

South and Central Palembang Sub-basins have shale hydrocarbon which are considerably promising in the Talang Akar and Lemat/Lahat Formations. Seismic interpretation results shows potential of shale hydrocarbon scattered in several areas around the Rukam-l, Kemang-l, Singa-l and Tepus'l wells. Generally, thermal modeling results indicate early maturity of oil on the value of  $R_o = 0.6\%$  at about 2000m depth (h), the oil generation value  $R_o = (0.7-0.9)\%$  at between (2200 £ h < 3100)m depth and the oil generation  $R_o$  values between  $(0.9-1.2)\%$  at a depth between (3100-3500m, Talang Akar and Lahat/Lemat Formations have a shallow marine depositional environment with Type II/III kerogen and lacustrine with Type III kerogen respectively. Based on advanced seismic data processing (seismic attributes and spectral decomposition) these areas are expected to have  $TOC > 2\%$ . The calculation (P-50) of potential unconventional oil and gas resources at Talang Akar and Lahat/Lemat Formations is estimated to be fairly large (up to 4200 MMBOE)

(Author)