

Comparison of laparoscopic nissen and toupet fundoplication using a propensity score matching analysis

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20472111&lokasi=lokal>

Abstrak

PURPOSE: Laparoscopic fundoplication (LF) has become a standard operative procedure for GERD-related diseases in Japan, although meta-analyses have mainly evaluated findings from Western countries. The propensity score matching method was used to compare and investigate the treatment outcomes of two fundoplication procedures (the Nissen and Toupet methods).

METHODS: Among 474 patients who underwent initial LF from December 1994 to April 2016, we extracted 401 cases (Nissen: 92 cases, Toupet: 309 cases), excluding 73 patients in whom follow-up was insufficient. We then matched 126 of these patients (63 per group).

RESULTS: The esophageal acid reflux time (%) was 12.2:2.8, being higher in the Nissen group than in the Toupet group ($p<0.001$). Regarding the surgical outcome, the amount of bleeding was higher in the Nissen group ($p=0.001$), and the number of hospitalization days following surgery was longer ($p=0.003$). Furthermore, a significantly rate of postoperative difficulty in swallowing (%) was observed in the Nissen group, at 13:0 ($p=0.004$). The recurrence rate (%) was 8:3, with no difference between the two groups ($p=0.243$).

CONCLUSIONS: Although there was no marked difference in the recurrence rate between the two procedures, postoperative dysphagia was observed at a higher frequency with the Nissen method than the Toupet method.