

Hubungan overweight dengan hipertensi remaja usia 15-17 tahun di Indonesia (Analisis lanjut data IFLS V tahun 2014) = The relationship between overweight with adolescent hypertension aged 15-17 years in Indonesia data (Analysis of IFLS V, 2014)

Erna Savitri, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20476629&lokasi=lokal>

Abstrak

Hipertensi remaja masih menjadi masalah bagi kesehatan karena dapat meningkatkan angka kesakitan dan kematian akibat penyakit kardiovaskular di masa dewasa. Salah satu faktor risiko hipertensi remaja adalah overweight. Terdapat peningkatan prevalensi overweight pada remaja umur 16-18 tahun di Indonesia berdasarkan Riskesdas tahun 2007-2013 sebesar 5,7% menjadi 6,7% Tujuan penelitian ini adalah mengetahui hubungan overweight dengan hipertensi remaja usia 15-17 tahun di Indonesia berdasarkan data IFLS V Tahun 2014.

Desain Penelitian ini adalah studi cross sectional dengan menggunakan data sekunder dari IFLS V Tahun 2014. Analisis data yang digunakan adalah Regresi Logistik. Hasil analisis menunjukkan remaja dengan overweight berpeluang 1,530 kali (95% CI; 1,080-2,166) dibandingkan remaja tidak overweight setelah dikendalikan oleh jenis kelamin. Remaja terutama remaja laki-laki dapat meningkatkan aktivitas fisik (jalan cepat dan lari) untuk mencegah overweight dan hipertensi.

<hr /><i>Adolescents hypertension is still become a problem for health, because it can increase the number of mortality and morbidity from cardiovascular disease in adulthood. Overweight has become one of the risk factors of it. There is increasing prevalence of overweight in adolescents of aged 16-18 in Indonesia. Based on Riskesdas, there is a growing of adolescents at the aged of 16-18 in Indonesia between 2007-2013, its around 5,7% and 6,7%. The aim of this study is to find out the relation of overweight and adolescents hypertension at the aged of 15-17 based on IFLS V data in 2014.</i>

The design of this study is using cross sectional secondary data of IFLS V 2014 with logistic regression are used for data analysis. The analysis showed that adolescents with overweight had a POR 1,530 times (95% CI 1.880-2.166) compared to un-overweight (normal-weight) adolescents after being control by gender. The boys have advantage to increasing of physical activity (by running or brisk walking) to prevent from overweight and hypertension.</i>