

Screening and preliminary optimizations for dihydroxyacetone production from glycerol by the gluconobacter and asaia isolates Found in Thailand / Issara Poljungreed, Siwarutt Boonyarattanakalin, Pattaraporn Yukphan

Issara Poljungreed, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20487620&lokasi=lokal>

Abstrak

ABSTRAK

This study aims to investigate the feasibility of using a minimal glycerol medium with inorganic salt supplementation for a low-cost dihydroxyacetone (DHA) production by bacteria. Acetic acid bacterial isolates of *Gluconobacter* and *Asaia*, which are found in the Northern, North-Eastern, Middle, Western, and Southern regions of Thailand, were evaluated for their ability to produce DHA using glycerol as a carbon source. During the qualitative screening, 66 isolates, including *Gluconobacter* (61 isolates) and *Asaia* (5 isolates), from a total of 486 isolates showed highly positive results by the Fehling test. The 5 isolates of *Asaia* gave low DHA production in the quantitative screening, whereas *Gluconobacter* isolates showed DHA production at low (0-5.70 g/L), medium (5.71-11.40 g/L), and high (11.41-16.89 g/L) levels. Preliminary culture medium optimizations for *G. frateurii* BCC 36199, a most promising microorganism for DHA production, were also carried out using a low-cost minimal glycerol medium supplemented with an inorganic salt. *G. frateurii* BCC 36199 produced 18.67 g/L of DHA with ysp of 95.44% (DHA moles/glycerol moles) at 30°C, 20 g/L of glycerol, and pH 4.5. The cultivation of *G. frateurii* BCC 36199 in the developed minimal glycerol medium is practical and can be further optimized in order to apply for industry.