

Luaran onkologi dan fungsional dari pasien metastasis tulang panjang dan axial Di RSUPN Dr Cipto Mangunkusumo Periode 2008-2018 = Oncological and functional outcome of patients with Long and Axial metastatic bone disease at Cipto Mangunkusumo Hospital on 2008 - 2018

Adisa Yusuf Reksoprodjo, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20494051&lokasi=lokal>

Abstrak

Metastatic Bone Disease (MBD) merupakan tempat penyebaran jauh terbanyak ketiga setelah paru dan liver. Hal ini menimbulkan morbiditas yang tidak sedikit dan pada akhirnya memengaruhi kualitas hidup dan kesintasan pasien. Penelitian ini menggunakan studi potong lintang di RSUPN Cipto Mangunkusumo dengan total sampling. Pasien yang terdiagnosis MBD selama periode 2008 - 2018 dilihat karakteristik, kesintasan, dan jika masih hidup, dilakukan penghitungan skor fungsional menggunakan kuesioner SF-36 dan MSTs. Terdapat 113 pasien MBD dengan rerata usia $54,34 \pm 11,09$, 69% perempuan, 24,8% tumor primer dari paru, 17,7% dari mammae, 16,8% dari tiroid. 55,8% lesi MBD terdapat pada ekstremitas dan 74,3% merupakan lesi soliter. 65,5% pasien tidak menjalani operasi, namun 78,8% mendapatkan bisfosfonat dan 51,3% mendapatkan radioterapi. Sebanyak 82,3% pasien sudah meninggal, sehingga terdapat 20 pasien yang masih hidup. SF-36 menunjukkan rentang median 40,0 - 100,0 dari 8 skala yang ada. MSTs ekstremitas atas rerata $45,55 \pm 24,46$ dan ekstremitas bawah median 26,67 (20,00 - 60,00). Analisis bivariat menunjukkan hubungan antara pembedahan dengan kesintasan ($P=0,034$). Analisis multivariat menunjukkan operasi memiliki peluang terhadap kesintasan yang lebih baik sebesar 2,8 kali (95%CI 1,1 - 7,6). Operasi memiliki hubungan yang bermakna terhadap kesintasan pasien MBD.

.....Metastatic Bone Disease (MBD) is the third distant sites after lungs and liver. This creates morbidity and affect patient s quality of life and survival. This study uses cross sectional design with total sampling at one tertiary referral center. MBD patient during 2008 - 2018 were evaluated for characteristics, treatment received, and survival rate. Survived patients were evaluated for functional score with SF-36 and MSTs. From 113 patients, with mean age of $54,34 \pm 11,09$, 69% were female, 24,8% were lung primary tumor, 17,7% from breast tumor, and 16,8% from thyroid tumor. 55,8% of the lesions were from extremity and 74,3% were solitary lesions. 65,5% patients did not get a surgery, 78,8% were given bisphosphonates, and 51,3% got a radiotherapy treatment. 82,3% patients were already died, so we got 20 patients that were still alive and being evaluated for the functional score. SF-36 shows median of 40,0 - 100,0 from 8 scales, and upper extremity MSTs results mean $45,55 \pm 24,46$, and lower extremity MSTs results median 26,67 (20,00 - 60,00). Bivariate analysis shows statistically significant association of surgery with survival ($P=0,034$). Multivariate analysis shows surgery has a 2,8 times higher chance of survival (95%CI 1,1 - 7,6). Surgery has a significant association with MBD patient survival.