

Gangguan keseimbangan tubuh pada pekerja tambang terpajan merkuri di kawasan pertambangan emas skala kecil (PESK) di Kecamatan Hutabargot Kabupaten Mandailing Natal Provinsi Sumatera Utara = Body balance disorder on mine workers exposed by mercury in the Small Scale Gold Mining Area (ASGM) in the sub-district of Hutabargot District Mandailing District, North Sumatera Province

Muhammad Rizhkal, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20494255&lokasi=lokal>

Abstrak

Pertambangan emas di kabupaten Mandailing Natal sudah ada sejak 2008. Tetapi semakin marak pada tahun 2010 di kecamatan Hutabargot dan di kecamatan Nagajuang pada November 2011. Tujuan dari penelitian ini adalah untuk menganalisis hubungan antara kadar merkuri pada rambut pada pekerja tambang terpajan merkuri dan karakteristik individu pekerja(usia, lama tinggal, status gizi dan konsumsi ikan) dengan gangguan keseimbangan tubuh. Penelitian ini menggunakan metode cross sectional. Data yang digunakan yaitu data primer dari kuesioner dan sekunder dari hasil uji laboratorium rambut. Sampel penelitian ini disesuaikan dengan sampel dari data sekunder yang menggunakan rumus Lemeshow sehingga didapatkan sampel 60 orang.

Hasil penelitian ini didapatkan tidak ada hubungan yang signifikan antara variabel independent (merkuri dalam rambut) dan variabel confounding terhadap variabel dependent (gangguan keseimbangan tubuh). Walaupun hasil penelitian tidak menunjukkan hubungan yang signifikan jika dilihat dari nilai OR variabel merkuri dalam rambut masih tergolong tinggi yaitu 6,0 dan setelah dikontrol variabel karakteristik individu OR merkuri dalam rambut turun menjadi 4,92.

<hr>

Gold mining in Mandailing Natal regency has been around since 2008. But it became more prevalent in 2010 in Hutabargot sub-district and in Nagajuang sub-district in November 2011. The purpose of this study was to analyze the relationship between mercury levels in hair in mercury exposed workers and individual characteristics of workers (age, length of stay, nutritional status and consumption of fish) with body balance disorders. This study uses a cross sectional method. The data used are primary data from questionnaires and secondary from the results of hair laboratory tests. The research sample was adjusted to the sample from secondary data using the Lemeshow formula so that sample of 60 people was obtained.

The results of this study found that there was no significant relationship between independent variables (mercury in hair) and confounding variables on the dependent variable (body balance disorder). Although the results of the study did not show a significant relationship when viewed from the OR variable value of mercury in the hair was still relatively high at 6.0 and after being contracted the individual characteristic variable OR mercury in the hair dropped to 4.92.