

Risk assesment of air pollution exposure (NO₂, SO₂, total suspended particulate, and particulate matter 10 micron) and smoking habits on the lung function of bus drivers in Palembang City

Elvi Sunarsih, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20497479&lokasi=lokal>

Abstrak

ABSTRACT

Nitrogen dioxide (NO₂), sulfur dioxide (SO₂), particulate matter (PM₁₀), and total suspended particulate (TSP) are the most common and harmful air pollutants to humans. In a short period, air pollution exposure at 5 ppm for 10 minutes in humans causes dyspnea, and when the level is increased to 800 ppm could cause 100% mortality in animals. This study was an analytical study with implementing a cross-sectional design and risk analysis. One hundred subjects were involved in this study. The results showed that the mean value of the non-cancer Hazard Index (HI) for real-time exposure was NO₂: 1.85; SO₂: 2.92; TSP: 7.09; and PM₁₀: 11.7 (HI value >1). Test for forced vital capacity lung capacity to non-cancer risk estimation of NO₂, SO₂, TSP, and PM₁₀ indicated that there was no significant relationship (p-value >0.05). The variable of smoking habit is the most dominant variable (odds ratio [OR] = 12.542) which affects respiratory disorders. The exposure to NO₂, SO₂, TSP, and PM₁₀ in Palembang City bus drivers is considered hazardous to the health of subjects without cancer.