

Uji antifeedant ekstrak kasar Timun Laut *Synaptula reticulata* (Semper, 1868) terhadap ikan karang di Perairan Pulau Pramuka, Taman Nasional Kepulauan Seribu, DKI Jakarta = Antifeedant Assay of Sea Cucumber *Synaptula reticulata* (Semper, 1868) Crude Extract on Reef Fishes at Pramuka Island Waters, Seribu Islands National Park, DKI Jakarta

Firli Rahman Hakim Fauzi, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20497693&lokasi=lokal>

Abstrak

Synaptula reticulata merupakan timun laut berdinding tubuh tipis dan memiliki warna kontras. Uji antifeedant ekstrak kasar *Synaptula reticulata* telah dilakukan pada tanggal 6-14 November di kedalaman 3-5 m Perairan Pulau Pramuka, Taman Nasional Kepulauan Seribu, DKI Jakarta. Sampel *Synaptula reticulata* berjumlah 171 individu dan dilarutkan dengan metanol. Persentase ekstrak kasar *Synaptula reticulata* yang didapat sebesar 8% dan memiliki konsentrasi fisiologis 40 mg/mL. Uji antifeedant dilakukan dengan membandingkan respon makan ikan karang terhadap pakan uji dan pakan kontrol. Pakan uji adalah ekstrak kasar *Synaptula reticulata* yang dicampur jelly dan pelet komersil.

Pakan kontrol adalah campuran jelly dan pelet komersil tanpa dicampurkan ekstrak kasar *Synaptula reticulata*. Jumlah pakan uji yang dimakan sebanyak 3%, sedangkan jumlah pakan kontrol yang dimakan sebanyak 63%. Hasil uji statistik Chi-kuadrat pada tingkat kepercayaan 99% menunjukkan bahwa pemberian ekstrak kasar *Synaptula reticulata* berhubungan dengan respon makan ikan karang. Keeratan hubungan tersebut kuat, berdasarkan uji korelasi cremer ($C = 0,63$) terutama terhadap ikan karang family Pomacentridae dan Laberidae.

Synaptula reticulata is sea cucumber that has thin body wall with contrasting color. Antifeedant activity assay from crude extract of *Synaptula reticulata* was conducted on 6th--10th November 2018 in 3--5 m of depth Pramuka Island water, Kepulauan Seribu National Park, DKI Jakarta. 171 individual *Synaptula reticulata* were collected and extracted using methanol. Crude extract percentage of *Synaptula reticulata* was 8% with a physiological concentration of 40 mg/mL. Antifeedant assay was done by comparing between coral reef fish feeding response to artificial test food and control food. Test food ware constitute of crude extract of *Synaptula reticulata*, jelly and pellet.

Control food contained jelly and pellet only. The amount of test food eaten as much as 3%, while the amount of control food eaten as much as 63%. Chi-Square analysis with confidence level of 0,01 showed that crude extract of *Synaptula reticulata* was correlated with feeding response of reef fishes. Cramer correlation test showed that crude extract *Synaptula reticulata* strongly related with feeding response of the treatments on reef fishes, with correlation value of 0,6.