

Pengaruh Edukasi dan Latihan Ambulasi terhadap Fall Efficacy dan Status Fungsional pada Pasien Pascapembedahan Open Reduction and Internal Fixation Ekstremitas Bawah = The Effect of Education and Ambulation Training on Fall Efficacy and Functional Status in Postsurgery Patients with Open Reduction and Internal Fixation of Lower Extremity

Faraniara, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20508862&lokasi=lokal>

Abstrak

ABSTRAK

Perasaan takut akan jatuh dan penurunan status fungsional sering dialami pada pasien fraktur ekstremitas bawah yang telah menjalani pembedahan. Edukasi yang tepat diikuti dengan latihan ambulasi pada pasien pascapembedahan diperlukan untuk dapat meningkatkan status fungsional dan keyakinan pasien untuk melakukan ambulasi dini. Penelitian ini bertujuan untuk mengetahui pengaruh edukasi dan latihan ambulasi terhadap fall efficacy dan status fungsional pada pasien pascapembedahan ORIF ekstremitas bawah. Desain penelitian ini adalah kuasi eksperimen dengan rancangan pre test and post test without control group. Besar sampel pada penelitian ini adalah sebanyak 33 responden dengan consecutive sampling. Hasil penelitian ini menunjukkan bahwa terdapat perbedaan fall efficacy dan status fungsional antara sebelum dan setelah dilakukan edukasi dan latihan ambulasi (p value $< 0,001$). Penelitian ini menyimpulkan terdapat perbedaan nilai fall efficacy dan status fungsional setelah diberikan intervensi edukasi dengan latihan ambulasi

<hr>

ABSTRACT

Fear of falling and decreasing functional status are often experienced in patients with lower limb fractures who have undergone surgery. Appropriate education followed by ambulation training in postsurgery patients is needed to be able to increase the functional status and confidence of patients to carry out early ambulation. This study aims to determine the effect of education and ambulation training on fall efficacy and functional status in postsurgery patients with lower extremity ORIF. The design of this study was a quasy experimental with a pre test and post test design without control group. The sample size in this study was 33 respondents with consecutive sampling. The results of this study indicate that there are differences in fall efficacy and functional status between before and after education and ambulation training (p value <0.001). This study concluded that there were differences in fall efficacy and functional status after being given educational intervention with ambulation training