

Efek irbesartan terhadap kadar malondialdehid jantung dan serum tikus model penyakit ginjal kronis = The effect of irbesartan on the levels of cardiac and serum malondialdehyde on rat models of chronic kidney disease

Kezia Alicia Theresia, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20510236&lokasi=lokal>

Abstrak

Penyakit Ginjal Kronis (PGK) merupakan kondisi dengan penurunan laju filtrasi glomerulus (LFG).

Penurunan LFG mengaktivasi Sistem-Renin-Angiotensin-Aldosteron (RAAS) dan menyebabkan penumpukan toksin uremik yang meningkatkan stres oksidatif. Irbesartan adalah obat yang berperan dalam inhibisi RAAS dan diduga memiliki peranan dalam menurunkan stres oksidatif.

Tujuan: Mengetahui efek antioksidan Irbesartan dan pengaruhnya terhadap stres oksidatif jantung dan serum tikus model PGK metode 5/6 nefrektomi melalui pengukuran kadar malondialdehid.

Metode: Penelitian ini merupakan penelitian eksperimental menggunakan jaringan jantung dan serum tikus tersimpan. Sampel terdiri atas 3 kelompok, yaitu sham (S, jantung dan serum: n=4), nefrektomi 5/6 (N, jantung: n=4; serum: n=3) dan nefrektomi 5/6 dengan pemberian Irbesartan 20 mg/kgBB/hari selama 4 minggu (N+I, jantung dan serum: n=4). Kemudian dilakukan pengukuran kadar malondialdehid melalui uji Thiobarbituric Acid Reactive Substance Assay (TBARS) pada jaringan tersimpan. Analisis statistik dilakukan dengan SPSS v25.0 menggunakan uji One-Way ANOVA.

Hasil: Tidak terdapat perbedaan bermakna secara statistik ($p > 0,05$) antara ketiga kelompok pada pemeriksaan kadar malondialdehid jantung dan serum tikus (jantung: $p = 0,060$; serum: $p = 0,162$).

Kesimpulan: Tidak terdapat perbedaan bermakna pada kadar malondialdehid jantung dan serum tikus model PGK metode 5/6 nefrektomi dengan dan tanpa pemberian Irbesartan.

.....Background: Chronic kidney disease (CKD) is a condition that triggers a decrease in glomerular filtration rate. The reduction can activate Renin-Angiotensin-Aldosterone System (RAAS) and leads to an accumulation of oxidative stress. Irbesartan is a drug that functions to inhibit RAAS and is thought to have an effect on lowering the oxidative stress.

Objectives: To understand irbesartan's antioxidant effects and its impact on oxidative stress in cardiac tissues and serum on rat models of 5/6 nephrectomy-induced CKD through the measurement of malondialdehyde levels.

Methods: This study is an experimental study utilizing stored heart and serum. The samples consists of three groups, which are Sham (S, heart and serum: n=4), 5/6 nephrectomy (N, heart: n=4; serum: n=3), and 5/6 nephrectomy administered with Irbesartan 20 mg/kgW/day for 4 weeks (N+I, heart and serum: n=4). Statistical analyses were done using SPSS v25.0 and examined using One-Way ANOVA.

Results: There were no significant results between the three groups based on the levels of heart and serum malondialdehyde (heart: $p = 0.060$; serum: $p = 0.162$).

Conclusion: There were no significant differences in the levels of heart and serum malondialdehyde on rat models of 5/6 nephrectomy-induced CKD with the administration of Irbesartan compared and without the administration of Irbesartan.