

## The differences of serum complements and anti-dsdna levels between renal and non-renal manifestations in systematic lupus erythematosus

Bagus Putu Putra Suryana, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20513119&lokasi=lokal>

---

### Abstrak

**Background.** Systemic lupus erythematosus (SLE) has diverse clinical manifestations, including renal and non-renal. Renal manifestation is related to significant morbidity and mortality. SLE is also characterized by serological aberrations, including levels of complement C3, C4 and anti-dsDNA, but the association of them with clinical manifestations including renal and non-renal is unclear. This study investigated the associations of C3, C4 and anti-dsDNA levels with renal and non-renal manifestations in SLE patients.

**Method.** A cross-sectional study was conducted in the Polyclinic of Rheumatology, Dr. Saiful Anwar Hospital Malang. A number of 43 subjects fulfilled the 1997 American College of Rheumatology criteria participated in this study, that consisted of 11 patients with renal manifestation and 32 patients with non-renal manifestations. Serum C3 and C4 levels were measured using immunoturbidimetry, and serum anti-dsDNA levels were measured using enzyme-linked immunosorbent assays (ELISA). The independent T-test was used to compare C3 levels and the Mann-Whitney U test was used to compare C4 and anti-dsDNA levels between groups.

**Result.** SLE with renal manifestation had significant lower levels of serum C3 compare to non-renal manifestations (mean  $\hat{\pm}$  SD: 71.27  $\hat{\pm}$  32.65 mg/dL and 94.47  $\hat{\pm}$  26.29 mg/dL respectively,  $p=0.022$ ).

SLE with renal manifestation also had significantly lower levels of serum C4 compare to non-renal manifestations (mean  $\hat{\pm}$  SD: 14.55  $\hat{\pm}$  8.20 mg/dL and 25.50  $\hat{\pm}$  11.05 mg/dL respectively,  $p=0.002$ ).

Conversely, SLE with renal manifestation had significantly higher levels of serum anti-dsDNA compare to non-renal manifestations (mean  $\hat{\pm}$  SD: 249.27  $\hat{\pm}$  240.34 IU/mL and 109.91  $\hat{\pm}$  166.11 IU/mL respectively,  $p=0.014$ ).

**Conclusion.** SLE patients with renal manifestation have significantly lower levels of serum C3 and C4 and a higher level of serum anti-dsDNA than SLE patients with non-renal manifestations.