

# Analisis praktik residensi keperawatan medikal bedah pada keperawatan medikal bedah pasien gangguan sistem kardiovaskular dengan pendekatan model konservasi myra estrin levine di Rumah Sakit Jantung dan Pembuluh Darah Harapan Kita Jakarta = The analysis of clinical residency practice of medical-surgical nursing on cardiovascular disorder patients by applying levine's conservation model approach at the Cardiovascular Center Harapan Kita Hospital, Jakarta

Gde Yasa Antarika, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20521228&lokasi=lokal>

---

## Abstrak

Kecemasan dan nyeri merupakan masalah paling umum pada pasca operasi bedah jantung terbuka. Kedua hal tersebut akan menimbulkan efek negatif seperti meningkatkan penggunaan analgetik, komplikasi paru, gangguan tidur, waktu penyembuhan luka, lama serta biaya perawatan rumah sakit. Pranayama alternate nostril breathing merupakan terapi non-farmakologi yang dapat menurunkan kecemasan dan nyeri pasca bedah jantung. Evidence based nursing practice (EBNP) bertujuan menilai pengaruh pranayama alternate nostril breathing terhadap kecemasan dan nyeri pada pasien pasca bedah jantung. Subjek penelitian sebanyak 44 orang dibagi menjadi kelompok intervensi sejumlah 22 orang dan kelompok kontrol 22 orang. Kelompok intervensi mendapat perlakuan pranayama alternate nostril breathing sedangkan kelompok kontrol berupa relaksasi napas dalam. Penilaian kecemasan menggunakan The State-Trait Anxiety Inventory (STAI) sedangkan penilaian nyeri menggunakan Visual Analog Scale (VAS) 0-100 mm. Uji statistik penelitian menggunakan analisis bivariat dengan uji paired t test dan independent t test. Hasil penelitian menunjukkan pranayama alternate nostril breathing dapat menurunkan intensitas kecemasan dan nyeri pada kelompok intervensi ( $p$ -value  $<0,05$ ) dibandingkan kelompok kontrol. Pranayama alternate nostril breathing merupakan intervensi non-farmakologi dalam menurunkan kecemasan dan nyeri pasien pasca bedah jantung yang bersifat sederhana, murah, efektif, efesien, dan fleksibel. Penelitian lanjutan perlu dilakukan dengan menggunakan sampel lebih besar serta instrumen penilaian kecemasan pasca operasi yang lebih spesifik.

.....Anxiety and pain are the most common problems after open heart surgery. Both of them causes negative effects such as increasing the use of analgesics, pulmonary complications, sleep disturbances, wound healing time, hospital length of stay and cost. Pranayama alternate nostril breathing is a non-pharmacological therapy that can reduce anxiety and pain in post-cardiac surgery patients. Evidence based nursing practice (EBNP) aims to assess the effect of Pranayama alternate nostril breathing on anxiety and pain in post-cardiac surgery. The subjects were 44 people who divided into an intervention group of 22 people and a control group of 22 people. The intervention group received Pranayama alternate nostril breathing while the control group was get deep breathing relaxation. The State-Trait Anxiety Inventory (STAI) was used for anxiety assessment while the Visual Analog Scale (VAS) for pain assesment. The statistic test used bivariate analysis with paired t test and independent t test. Pranayama alternate nostril breathing could reduce the intensity of anxiety and pain in the intervention group ( $p$ -value  $<0.05$ ) compared to the control group. Pranayama alternate nostril breathing is a non-pharmacological intervention for reducing anxiety and pain in post-cardiac surgery patients that is simple, inexpensive, effective, efficient, and flexible. Further research needs to be done using a larger sample and a more specific postoperative

anxiety assessment instrument.