

Analisis asuhan keperawatan keluarga dengan gizi kurang dan penerapan intervensi pemberian makanan selingan bergizi sebagai upaya pencegahan tuberkulosis = Analysis of nursing care for families with malnutrition and application of interventions for providing nutritious interludes as an effort for tuberculosis prevention

Amyra Luthfia Mumpuni, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20522710&lokasi=lokal>

---

#### Abstrak

Gizi kurang adalah salah satu dari masalah kesehatan di area perkotaan yang sering dijumpai akibat dari faktor sosial ekonomi maupun pengetahuan Ibu. Masalah kesehatan tersebut merupakan salah satu faktor risiko penyakit tuberkulosis (TBC). Karya ilmiah ini menggambarkan asuhan keperawatan pada keluarga Bapak T dengan masalah ketidakseimbangan nutrisi kurang dari kebutuhan tubuh. Implementasi yang telah dilakukan terdiri dari implementasi yang bersifat kognitif, afektif, dan psikomotor dengan pendekatan lima tugas kesehatan keluarga. Intervensi unggulan berupa pemberian makanan selingan bergizi selama 3 minggu luring dan 1 minggu daring. Hasil evaluasi menunjukkan terjadinya peningkatan berat badan balita sebanyak 700 gram. Sehingga, intervensi pemberian makanan selingan bergizi dan peningkatan pengetahuan Ibu perlu ditingkatkan kembali agar terjadi peningkatan berat badan yang lebih signifikan sebagai upaya mencegah terjadinya TBC.

.....Malnutrition is one of the health problems in urban areas that is often encountered due to socio-economic factors and mother's knowledge. This health problem is one of the risk factors for tuberculosis (TB). This scientific work describes nursing care for Mr. T's family with the problem of nutritional imbalance less than the body's needs. The implementation that has been carried out consists of implementations that are cognitive, affective, and psychomotor with a five-task family health approach. The leading intervention is providing nutritious snacks for 3 weeks offline and 1 week online. The results of the evaluation showed an increase in the body weight of toddlers as much as 700 grams. Thus, the intervention of providing nutritious snacks and increasing mother's knowledge needs to be increased again so that there is a more significant increase in body weight as an effort to prevent the occurrence of tuberculosis.