

Hubungan Status Gizi Dengan Rasio Netrofil Limfosit Pada Pasien Kanker Paru di RSUP Persahabatan = The Relationship Between Nutritional Status And Neutrophil Lymphocyte Ratio in Patient with Lung Cancer at RSUP Persahabatan

Lidya Anissa, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920521776&lokasi=lokal>

Abstrak

Pada penderita kanker paru terjadi inflamasi sistemik dan dapat dilihat dengan peningkatan rasio netrofil limfosit di mana pemeriksaan ini lazim dilakukan di Rumah Sakit. Inflamasi sistemik dapat menyebabkan anoreksi sehingga asupan pada penderita kanker paru menurun dan memengaruhi status gizinya. Kejadian malnutrisi yang tinggi pada pasien paru dapat berakibat lamanya rawat inap, turunnya kualitas hidup, dan dapat memengaruhi keberhasilan terapi kanker sehingga terapi nutrisi yang cepat dan tepat sangat perlu dilakukan. Salah satu diagnostik status gizi pada penderita kanker yaitu dengan menggunakan kriteria ASPEN yang terdiri dari penurunan asupan, penurunan berat badan, penurunan massa otot dan massa lemak subkutan, akumulasi cairan general atau lokal, dan kapasitas fungsional. Dikatakan malnutrisi jika terdapat dua dari enam kriteria tersebut. Penelitian ini merupakan studi potong lintang yang bertujuan untuk mengetahui hubungan status gizi dengan rasio netrofil limfosit pada pasien kanker paru di RSUP Persahabatan. Data diambil dari wawancara, pemeriksaan fisis, pemeriksaan laboratorium, dan rekam medis pasien poliklinik onkologi RSUP Persahabatan ($n = 52$). Pada penelitian ini subjek sebagian besar berjenis laki-laki (61,5%), rentang usia terbanyak antara 50-60 tahun (38,5%), memiliki riwayat merokok (55,8%) dengan indeks Brinkman berat (30,8%). Lebih dari 50% subjek dengan asupan energi dan protein dibawah rekomendasi asupan untuk pasien kanker. Sebagian besar subjek penelitian berisiko malnutrisi atau malnutrisi sedang (38,5%) dan sebanyak 67,3% mengalami malnutrisi. Pada penelitian ini subjek dengan nilai rasio netrofil limfosit tinggi sebanyak 38,5% dan rendah sebanyak 61,5%. Tidak terdapat hubungan antara status gizi dengan rasio netrofil limfosit pada penelitian ini ($p = 0,35$).

.....Systemic inflammation in patients with lung cancer can be seen by the increase in the neutrophil lymphocyte ratio where these examinations are common in hospitals. Systemic inflammation can cause anorexia, with the result that nutrition intake of lung cancer patients decreases and affects their nutritional status. High incidence of malnutrition in lung cancer patients can result in length of stay, decreased quality of life, and can affect the treatment of cancer therapy, therefore prompt and appropriate nutritional therapy is essential. One of the diagnostics of nutritional status for lung cancer patients is by using the ASPEN criteria which consist of decreased nutritional intake, weight loss, decreased muscle mass and subcutaneous fat mass, general or local fluid accumulation, and functional capacity. Malnutrition can be seen if there are two of the six criteria. This study is a cross-sectional study which aimed to determine the association between nutritional status and the ratio of lymphocyte neutrophils in lung cancer patients at Persahabatan Hospital. Data were taken from interviews, physical examinations, laboratory analysis, and patients medical records in the oncology polyclinic of Persahabatan Hospital ($n = 52$). The subjects of the study were mostly male (61.5%), the largest age range was between 50-60 years (38.5%), had a history of smoking (55.8%) with a severe Brinkman index (30.8%). More than 50% of the subjects with energy and protein intake were below

the recommended intake for cancer patients. Most of the study subjects were at risk of malnutrition or moderate malnutrition (38.5%) and 67.3% of them were experiencing malnutrition. The subjects with the highest neutrophil lymphocyte ratio value were 38.5% and the lowest value were 61.5%. Overall, there was no relationship between nutritional status and the ratio of neutrophil to lymphocytes in this study ($p = 0.35$).