

Analisis Karakteristik Sosiodemografi, Faktor Perilaku, dan Faktor Lingkungan Rumah Terhadap Kejadian Tuberkulosis Paru pada Penduduk Berusia 15 Tahun di Indonesia (Analisis Data Riskesdas 2018) = "Analysis of Sociodemographic Characteristics, Behavioral Factors, and Home Environment Factors on the Incidence of Pulmonary Tuberculosis in Populations Aged 15 Years in Indonesia (2018 Riskesdas Data Analysis)"

Risa Paradilla Utami, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920525054&lokasi=lokal>

Abstrak

Latar belakang: Indonesia merupakan negara dengan beban tuberkulosis tertinggi kedua di dunia setelah India dengan perkiraan morbiditas sebanyak 969.000 dan mortalitas mencapai 144.000 orang pada tahun 2021. Tujuan: Penelitian ini bertujuan untuk mengetahui hubungan antara karakteristik sosiodemografi, faktor perilaku, dan faktor lingkungan rumah terhadap kejadian tuberkulosis paru pada penduduk usia 15 tahun di Indonesia. Metode: Metode yang digunakan dalam penelitian ini adalah cross-sectional. Hasil: Variabel yang ditemukan berhubungan dengan kejadian TB paru adalah usia (OR = 2,107, 95% CI = 1,919-2,314), jenis kelamin (OR = 1,469, 95% CI = 1,371-1,575), status kawin (OR = 1,206, 95% CI = 1,117-1,303), tingkat pendidikan (OR = 1,795, 95% CI = 1,655-1,946), riwayat merokok (OR = 1,194, 95% CI = 1,113-1,281), kebiasaan membuka jendela rumah (OR = 1,160, 95% CI = 1,080-1,246), kondisi ventilasi (OR = 1,266, 95% CI = 1,178-1,360), kondisi pencahayaan (OR = 1,330, 95% CI = 1,241-1,426), jumlah anggota rumah tangga (OR = 1,131, 95% CI = 1,044-1,221), dan daerah tempat tinggal (OR = 1,213, 95% CI = 1,130-1,301). Riwayat konsumsi minuman beralkohol ditemukan sebagai faktor protektif. Kesimpulan: Terdapat hubungan antara karakteristik sosiodemografi, faktor perilaku, dan faktor lingkungan rumah dengan kejadian tuberkulosis paru pada penduduk usia 15 tahun di Indonesia.

.....Background: Indonesia is a country with the second highest tuberculosis burden in the world after India with an estimated morbidity of 969,000 and mortality reaching 144,000 people in 2021. Objective: This study aims to determine the relationship between sociodemographic characteristics, behavioral factors, and home environmental factors on incidence pulmonary tuberculosis in population aged 15 years in Indonesia. Methods: The method used in this study was cross-sectional. Results: The variables found to be associated with the incidence of pulmonary TB were age (OR = 2.107, 95% CI = 1.919-2.314), gender (OR = 1.469, 95% CI = 1.371-1.575), marital status (OR = 1.206, 95 % CI = 1.117-1.303), education level (OR = 1.795, 95% CI = 1.655-1.946), smoking history (OR = 1.194, 95% CI = 1.113-1.281), habit of opening windows (OR = 1.160, 95 % CI = 1.080-1.246), ventilation conditions (OR = 1.266, 95% CI = 1.178-1.360), lighting conditions (OR = 1.330, 95% CI = 1.241-1.426), number of household members (OR = 1.131, 95 % CI = 1.044-1.221), and area of residence (OR = 1.213, 95% CI = 1.130-1.301). History of alcohol consumption was found to be a protective factor. Conclusion: There is a relationship between sociodemographic characteristics, behavioral factors, and home environment factors with the incidence of pulmonary tuberculosis in people aged 15 years in Indonesia.