

Efektivitas Pemberian Intervensi Keperawatan Range of Motion (ROM) dan Terapi Genggam Bola Terhadap Peningkatan Kekuatan Otot pada Keluarga yang Mengalami Stroke = The Effectiveness of Providing Range of Motion (ROM) Nursing Interventions and Ball Grasp Therapy on Increasing Muscle Strength in Families Experiencing Stroke

Showmi Ati Aisyah, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920537064&lokasi=lokal>

Abstrak

Seseorang yang terkena stroke akan mengalami gangguan mobilitas, kelumpuhan atau kelemahan satu sisi (Hemiparesis) yang akan mengakibatkan menurunnya kemampuan merawat diri. Stroke menyerang pada usia dewasa dan penyebab stroke yakni hipertensi dan kolesterol tinggi. Metode yang digunakan yaitu asuhan keperawatan mulai pengkajian hingga evaluasi penderita stroke yang mengalami keluhan kaku sendi pada ekstremitas kanan yakni kaki dan tangan. Tujuan penelitian ini mengetahui efektifitas ROM dan terapi genggam bola. Hasil implementasi ini berlangsung selama 14 kali dalam waktu 45 menit selama 11 kali pertemuan pada ROM terjadi peningkatan kekuatan otot senilai 2 pada ekstremitas bawah terjadi peningkatan kekuatan senilai 3 selanjutnya untuk intervensi terapi genggam bola mengalami peningkatan otot pada ekstremitas atas di pergelangan tangan senilai 1 dan pada jari-jari mengalami peningkatan 2 selama 11 kali Latihan terapi genggam bola.

.....A person who has had a stroke will experience impaired mobility, paralysis or one-sided weakness (Hemiparesis) which will result in decreased ability to care for themselves. The method used is nursing care from assessment to evaluation of stroke patients who experience complaints of stiff joints in the right extremities, namely the feet and hands. The purpose of this study was to determine the effectiveness of ROM and ball-grasping therapy. The results of this implementation lasted 14 times within 45 minutes for 11 meetings at ROM there was an increase in muscle strength worth 2 in the lower extremities, and there was an increase in strength worth 3 then for the ball grasp therapy intervention experienced an increase in muscle in the upper extremities in the wrist worth 1 and in the fingers experienced an increase of 2 during 11 times of ball grasp therapy exercises.