

# Pengaruh Terapi Akupresur Terhadap Perbaikan Klinis Ulkus Diabetik = Effect of Acupressure to the Clinical Improvement of Diabetic Ulcer

Tommy Indra, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=9999920541245&lokasi=lokal>

---

## Abstrak

Penderita diabetes melitus jangka panjang dapat menyebabkan komplikasi mikrovaskular dan makrovaskular yang mengakibatkan peredaran darah arteri yang menyempit sehingga mengurangi aliran darah ke tungkai bawah. Hal ini dapat mengakibatkan sirkulasi oksigen yang buruk sehingga berdampak meningkatkan risiko terjadinya ulkus diabetik. Ulkus diabetik merupakan penyebab morbiditas dan mortalitas yang signifikan, jika tidak ditangani dan diobati secara tepat waktu akan menyebabkan amputasi. Hal ini perlu dilakukan tindakan pencegahan dan pengelolaan ulkus diabetik yang baik untuk mengatasi tersebut. Salah satu tindakan komplementer yaitu terapi akupresur, dapat dimanfaatkan pada penderita ulkus diabetik. Tujuan dari penelitian ini yaitu mengidentifikasi pengaruh akupresur terhadap perbaikan klinis ulkus diabetik. Sampel penelitian ini berjumlah 26 responden yang dibagi menjadi 13 responden kelompok intervensi dan 13 responden kelompok kontrol. Metode penelitian ini adalah *Quasi Eksperimen* dengan desain penelitian *Pretest and Posttest With Control Group Designs*. Hasil penelitian menunjukkan bahwa ada perbedaan yang cukup signifikan rerata selisih skor ulkus diabetik antara kelompok intervensi dengan kelompok kontrol dengan *p value* 0,047 (*p value* <0,05). Hasil uji variabel *confounding* didapat data bahwa lama menyandang DM dan riwayat merokok, kadar glukosa darah sewaktu dan tekanan darah tidak berhubungan dengan perbaikan klinis ulkus diabetik (*p value* > 0,05). Terapi akupresur diharapkan dapat menjadi standar terapi komplementer dalam pelayanan kesehatan dirumah sakit maupun pelayanan di klinik keperawatan luka sebagai terapi tambahan pada manajemen perawatan luka pada pasien ulkus diabetik.

Long-term diabetes mellitus patients can cause microvascular and macrovascular complications resulting in narrowed arterial blood circulation thereby reducing blood flow to the lower limbs. This can result in poor oxygen circulation thus increasing the risk of developing diabetic ulcers. Diabetic Ulcer is a significant cause of morbidity and mortality, which in case of not taken care or treated timely, it will lead to amputation. This requires good prevention and management of diabetic ulcer to handle such matter. One of the complimentary actions to help diabetic ulcer patients is acupressure. Objective of this research is to identify the impacts of acupressure to the clinical improvement of diabetic ulcer. Samples of this research contain 26 respondents, which come from 13 respondents of intervention group and 13 respondents of control group. Method of this research is *Quasi-Experiment with control group pretest-posttest design*. The research result indicates that acupressure with modern wound care decreases the score of diabetic ulcer improvement compared to standard therapy only (modern wound care) with *p value* 0,047. Results of the confounding variable test show that the terms of DM disease, smoking history, blood glucose levels and blood pressure do not relate to the clinical improvement of diabetic ulcer (*p value* > 0.05). Acupressure is expected may become a standard complimentary therapy of medical treatment in hospitals or wound care clinics as an additional therapy in diabetic wound care.